

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

APPLICATION FOR  
UNITED STATES LETTERS PATENT

This application claims benefit of co-pending U.S. Patent Application Serial No. 60/264,649 filed January 26, 2001, entitled "Mosquito Olfactory Genes, Polypeptides, and Methods of Use Thereof" which is hereby incorporated by reference. Be it known that I, Laurence J Zwiebel, a citizen of the United States, residing at 2512 Sunset Place, Nashville, TN 37212; have invented a new and useful "Mosquito Olfactory Genes, Polypeptides, and Methods of Use Thereof".

GOVERNMENT SUPPORT CLAUSE

This invention was made with federal grant money under NIH grant 1 R01 DC04692-01 and NSF grant 0075338. The United States Government has certain rights in this invention.

A Portion of the disclosure of this patent document contains material which is subject to copyright protection. The copyright owner has no objection to the facsimile reproduction by anyone of the patent document or the patent disclosure, as it appears

in the Patent and Trademark Office patent file or records, but otherwise reserves all copyright rights whatsoever.

## FIELD OF THE INVENTION

5       The present invention relates generally to the field of host identification by insects. Specifically, the present invention relates to the identification and cloning of genes related to mosquito olfaction, identification and purification of polypeptides thereof, and methods of use thereof.

## BACKGROUND OF THE INVENTION

10       The ability of an insect to respond to chemical stimuli is necessary for the insect to reproduce, mate, and feed. For example, insects respond to certain chemical stimuli by moving up a chemical gradient to identify and target a host. Mosquitoes, in particular, are believed to use olfaction to identify and target sources of bloodmeal for  
15       reproductive purposes. This behavior contributes to the spread of diseases in humans, such as malaria, encephalitis, and dengue fever; as well as, animal and livestock disease.

20       Olfaction plays a critical role in insect behaviors among agricultural pests and disease vectors. Hildebrand, et al., 1997, Annu. Rev. Neurosci, 20:595-631. In *Drosophila melanogaster* (the common fruit fly), the olfactory system functions through a rapid cycling between an on and off state of certain regulatory molecules.

10056409-01409  
204370-049500T

The olfactory signal transduction cascade is “turned on” by ligand-based activation of an odorant receptor and transduction of the signal by G-protein coupled second messenger pathways Boekhoff *et al.*, 1994, J. Neurosci, 14:3304-9. The “on signal” is rapidly and substantially terminated in the *Drosophila* system through the modification of the odorant receptor such that the G-protein coupled second messenger pathway is deactivated. Dohlman *et al.*, 1991, Annual Review of Biochemistry, 60:653-88. Olfactory transduction is provided by second messenger pathways of G protein-coupled receptors. Reed, R., 1992, Neuron 8:205-209; Bloekhoff, *et al*, 1994, Neurosci 14:3304-3309.

The structural and functional characteristics of the mosquito olfactory system has not been characterized to date. Given the importance of the controlling this pest and disease vector, what is needed is the identification and characterization of the genes and polypeptides that function for mosquito olfaction and methods of use thereof for mosquito management.

## SUMMARY OF THE INVENTION

The present invention provides, in part, eight novel mosquito polypeptides and nucleic acids encoding the polypeptides (collectively referred to herein as “mosquito olfaction molecules”). Seven of the polypeptides are novel mosquito odorant receptors and the eighth is a novel mosquito arrestin molecule (see Figure 8). The odorant receptor molecules are discovered to function in a ligand-induced signal transduction

pathway for the activation of mosquito olfaction. The mosquito arrestin molecule is discovered to function to inhibit the activated signal transduction cascade. Thus, the odorant receptors can be viewed as parts of an "on switch" or an "on signal" and the arrestin molecule can be viewed as an "off switch" or an "off signal" for the odorant  
5 detection system of the mosquito. The present invention is not bound by theory or mechanism.

The present invention also provides, in part, a system for disrupting the mosquito olfactory system by disrupting, inhibiting, or otherwise interfering with the function of the off switch for mosquito olfaction. Such interference is contemplated to inhibit or degrade the ability of the mosquito to appropriately  
10 respond to chemical clues in the environment used by the mosquito for host identification and targeting. For, example, if the signal cascade cannot be terminated or inhibited, then the mosquito is impaired in following a chemical gradient to a host through sampling of the frequency of ligand-induced activation of  
15 the olfaction signal cascade. In this example, the chemical concentration of the odorant is expected to increase with decreasing distance to the target. Thus, receptor activation is expected to increase with decreasing distance to the target. It is a discovery of the present invention, that factors that inhibit the on and off cycling of the mosquito olfactory signal cascade through inhibition of signal  
20 deactivation are useful for the control of mosquitoes. Test agents used in a method for identifying mosquito olfaction molecule binding compounds would include, but

are not limited to: chemicals, proteins, peptides, organic compounds and lipids. Such factors that inhibit signal deactivation may be peptides and chemicals. Several classes of chemicals that would be selected as targets are the carboxylic acids and steroids that are components of human sweat. Cork, A. (1996). Olfactory sensing is the basis of host location by mosquitoes and other hematophagous Diptera. In Olfaction in Mosquito-Host Interactions, G. R. B. a. G. Cardew, ed. (Chichester, New York, Brisbane, Toronto , Singapor: John Wiley & Sons), pp. 71-84. Furthermore, certain aspects of the present invention are contemplated to be effective for insects in general.

Methods are presented for identifying compounds that interfere with the operation of the mosquito olfactory system resulting in an over stimulation of olfactory signaling. One consequence of interfering with the mosquito olfactory system is that the mosquito has a diminished ability to home in on sources of bloodmeal. Additionally, interfering with mosquito insect olfactory systems will inhibit mating and feeding having a significant impact on mosquito populations and is helpful, for example, in nuisance and disease vector control for humans and livestock. Interfering with non-mosquito insect olfaction will similarly have a positive impact in control of other insect populations including for the protection of crops, such as: wheat, corn, rice, cotton, and soybeans. Thus, certain aspects of the present invention provide screening assays for the identification of compositions that will reduce the ability of mosquitoes to locate sources of bloodmeal, such as humans and other mammals,

including livestock (cattle, pigs, horses, sheep, etc.), show animals (horses, pigs, sheep, dogs, cats, etc.), and pets (dogs, cats, horses, etc). Certain aspects of the present invention provide a screening assay for the production of "mosquito olfaction molecules."

5 One aspect of the present invention provides an isolated DNA comprising a nucleotide sequence that encodes arrestin 1 polypeptide (e.g., SEQ ID NO: 2). In certain embodiments, arrestin 1 nucleotide sequence comprises a DNA molecule that hybridizes under stringent conditions to a DNA having a nucleotide sequence consisting of SEQ ID NO: 1, or the complement of SEQ ID NO: 1. Preferably the  
10 isolated DNA encodes naturally-occurring *Anopheles gambiae* arrestin 1 polypeptides. In certain embodiments, the nucleotide sequence may be that of SEQ ID NO: 1. In alternate embodiments, the nucleotide sequence may encode a fragment of SEQ ID NO: 2 at least 20 residues in length. One of ordinary skill in the art knows that a polypeptide fragment having a length of 20 residues is capable of functioning as an  
15 immunogen. In certain embodiments, the nucleotide sequence may encode a polypeptide having a conservatively modified amino acid sequence of SEQ ID NO: 2. In certain embodiments, the isolated polynucleotide comprises a complement to a sequence that encodes a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 2, and conservatively modified SEQ ID NO: 2.  
20 In alternate embodiments, the nucleotide sequence may be that of degenerate variants of above-mentioned sequences. The invention also includes operably linking one or

more expression control sequences to any of the above-mentioned nucleotide sequences. The invention also includes a cell comprising any of the above-mentioned nucleotide sequences operably linked to one or more expression control sequences.

The present invention also provides an isolated DNA comprising a nucleotide sequence that encodes odorant receptor 1 polypeptide (e.g., SEQ ID NO: 4). In certain embodiments, odorant receptor 1 nucleotide sequence comprises a DNA molecule that hybridizes under stringent conditions to a DNA having a nucleotide sequence consisting of SEQ ID NO: 3, or the complement of SEQ ID NO: 3. Preferably the isolated DNA encodes naturally-occurring *Anopheles gambiae* odorant receptor 1 polypeptides. In certain embodiments, the nucleotide sequence may be that of SEQ ID NO: 3. In alternate embodiments, the nucleotide sequence may encode a fragment of SEQ ID NO: 4 at least 20 residues in length. One of ordinary skill in the art knows that a polypeptide fragment having a length of 20 residues is capable of functioning as an immunogen. In certain embodiments, the nucleotide sequence may encode a polypeptide having a conservatively modified amino acid sequence of SEQ ID NO: 4. In certain embodiments, the isolated polynucleotide comprises a complement to a sequence that encodes a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 4, and conservatively modified SEQ ID NO: 4. In other alternate embodiments, the nucleotide sequence may be that of degenerate variants of above-mentioned sequences. The invention also includes operably linking one or more expression control sequences to any of the above-mentioned nucleotide

sequences. The invention also includes a cell comprising any of the above-mentioned nucleotide sequences operably linked to one or more expression control sequences.

The present invention provides an isolated DNA comprising a nucleotide sequence that encodes odorant receptor 2 polypeptide (e.g., SEQ ID NO: 6). In certain  
5 embodiments, odorant receptor 2 nucleotide sequence comprises a DNA molecule that hybridizes under stringent conditions to a DNA having a nucleotide sequence consisting of SEQ ID NO: 5, or the complement of SEQ ID NO: 5. Preferably the isolated DNA encodes naturally-occurring *Anopheles gambiae* odorant receptor 2 polypeptides. In certain embodiments, the nucleotide sequence may be that of SEQ ID  
10 NO: 5. In alternate embodiments, the nucleotide sequence may encode a fragment of SEQ ID NO: 6 at least 20 residues in length. One of ordinary skill in the art knows that a polypeptide fragment having a length of 20 residues is capable of functioning as an immunogen. In certain embodiments, the nucleotide sequence may encode a polypeptide having a conservatively modified amino acid sequence of SEQ ID NO: 6.  
15 In certain embodiments, the isolated polynucleotide comprises a complement to a sequence that encodes a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 6, and conservatively modified SEQ ID NO: 6. In other alternate embodiments, the nucleotide sequence may be that of degenerate variants of above-mentioned sequences. The invention also includes operably linking  
20 one or more expression control sequences to any of the above-mentioned nucleotide



sequences. The invention also includes a cell comprising any of the above-mentioned nucleotide sequences operably linked to one or more expression control sequences.

The present invention also provides an isolated DNA comprising a nucleotide sequence that encodes odorant receptor 3 polypeptide (e.g., SEQ ID NO: 8). In certain  
5 embodiments, odorant receptor 3 nucleotide sequence comprises a DNA molecule that hybridizes under stringent conditions to a DNA having a nucleotide sequence consisting of SEQ ID NO: 7, or the complement of SEQ ID NO: 7. Preferably the isolated DNA encodes naturally-occurring *Anopheles gambiae* odorant receptor 3 polypeptides. In certain embodiments, the nucleotide sequence may be that of SEQ ID  
10 NO: 7. In alternate embodiments, the nucleotide sequence may encode a fragment of SEQ ID NO: 8 at least 20 residues in length. One of ordinary skill in the art knows that a polypeptide fragment having a length of 20 residues is capable of functioning as an immunogen. In certain embodiments, the nucleotide sequence may encode a polypeptide having a conservatively modified amino acid sequence of SEQ ID NO: 8.

15 In certain embodiments, the isolated polynucleotide comprises a complement to a sequence that encodes a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 8, and conservatively modified SEQ ID NO: 8. In other alternate embodiments, the nucleotide sequence may be that of degenerate variants of above-mentioned sequences. The invention also includes operably linking  
20 one or more expression control sequences to any of the above-mentioned nucleotide

sequences. The invention also includes a cell comprising any of the above-mentioned nucleotide sequences operably linked to one or more expression control sequences.

The present invention also provides an isolated DNA comprising a nucleotide sequence that encodes odorant receptor 4 polypeptide (*e.g.*, SEQ ID NO: 14). In certain  
5 embodiments, odorant receptor 4 nucleotide sequence comprises a DNA molecule that hybridizes under stringent conditions to a DNA having a nucleotide sequence consisting of SEQ ID NO: 13, or the complement of SEQ ID NO: 13. Preferably the isolated DNA encodes naturally-occurring *Anopheles gambiae* odorant receptor 4 polypeptides. In certain embodiments, the nucleotide sequence may be that of SEQ ID  
10 NO: 13. In alternate embodiments, the nucleotide sequence may encode a fragment of SEQ ID NO: 14 at least 20 residues in length. One of ordinary skill in the art knows that a polypeptide fragment having a length of 20 residues is capable of functioning as an immunogen. In certain embodiments, the nucleotide sequence may encode a polypeptide having a conservatively modified amino acid sequence of SEQ ID NO: 14.  
15 In certain embodiments, the isolated polynucleotide comprises a complement to a sequence that encodes a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 14, and conservatively modified SEQ ID NO: 14. In other alternate embodiments, the nucleotide sequence may be that of degenerate variants of above-mentioned sequences. The invention also includes  
20 operably linking one or more expression control sequences to any of the above-mentioned nucleotide sequences. The invention also includes a cell comprising any of

the above-mentioned nucleotide sequences operably linked to one or more expression control sequences.

The present invention also provides an isolated DNA comprising a nucleotide sequence that encodes odorant receptor 5 polypeptide (*e.g.*, SEQ ID NO: 16). In certain embodiments, odorant receptor 5 nucleotide sequence comprises a DNA molecule that hybridizes under stringent conditions to a DNA having a nucleotide sequence consisting of SEQ ID NO: 15, or the complement of SEQ ID NO: 15. Preferably the isolated DNA encodes naturally-occurring *Anopheles gambiae* odorant receptor 5 polypeptides. In certain embodiments, the nucleotide sequence may be that of SEQ ID NO: 15. In alternate embodiments, the nucleotide sequence may encode a fragment of SEQ ID NO: 16 at least 20 residues in length. One of ordinary skill in the art knows that a polypeptide fragment having a length of 20 residues is capable of functioning as an immunogen. In certain embodiments, the nucleotide sequence may encode a polypeptide having a conservatively modified amino acid sequence of SEQ ID NO: 16. In certain embodiments, the isolated polynucleotide comprises a complement to a sequence that encodes a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 16, and conservatively modified SEQ ID NO: 16. In other alternate embodiments, the nucleotide sequence may be that of degenerate variants of above-mentioned sequences. The invention also includes operably linking one or more expression control sequences to any of the above-mentioned nucleotide sequences. The invention also includes a cell comprising any of

the above-mentioned nucleotide sequences operably linked to one or more expression control sequences.

The present invention also provides an isolated DNA comprising a nucleotide sequence that encodes odorant receptor 6 polypeptide (*e.g.*, SEQ ID NO: 18). In certain embodiments, odorant receptor 6 nucleotide sequence comprises a DNA molecule that hybridizes under stringent conditions to a DNA having a nucleotide sequence consisting of SEQ ID NO: 17, or the complement of SEQ ID NO: 17. Preferably the isolated DNA encodes naturally-occurring *Anopheles gambiae* odorant receptor 6 polypeptides. In certain embodiments, the nucleotide sequence may be that of SEQ ID NO: 17. In alternate embodiments, the nucleotide sequence may encode a fragment of SEQ ID NO: 18 at least 20 residues in length. One of ordinary skill in the art knows that a polypeptide fragment having a length of 20 residues is capable of functioning as an immunogen. In certain embodiments, the nucleotide sequence may encode a polypeptide having a conservatively modified amino acid sequence of SEQ ID NO: 18. In certain embodiments, the isolated polynucleotide comprises a complement to a sequence that encodes a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 18, and conservatively modified SEQ ID NO: 18. In other alternate embodiments, the nucleotide sequence may be that of degenerate variants of above-mentioned sequences. The invention also includes operably linking one or more expression control sequences to any of the above-mentioned nucleotide sequences. The invention also includes a cell comprising any of

the above-mentioned nucleotide sequences operably linked to one or more expression control sequences.

The present invention also provides an isolated DNA comprising a nucleotide sequence that encodes odorant receptor 7 polypeptide (*e.g.*, SEQ ID NO: 20). In certain  
5 embodiments, odorant receptor 7 nucleotide sequence comprises a DNA molecule that hybridizes under stringent conditions to a DNA having a nucleotide sequence consisting of SEQ ID NO: 19, or the complement of SEQ ID NO: 19. Preferably the isolated DNA encodes naturally-occurring *Anopheles gambiae* odorant receptor 7 polypeptides. In certain embodiments, the nucleotide sequence may be that of SEQ ID  
10 NO: 19. In alternate embodiments, the nucleotide sequence may encode a fragment of SEQ ID NO: 20 at least 20 residues in length. One of ordinary skill in the art knows that a polypeptide fragment having a length of 20 residues is capable of functioning as an immunogen. In certain embodiments, the nucleotide sequence may encode a polypeptide having a conservatively modified amino acid sequence of SEQ ID NO: 20.  
15 In certain embodiments, the isolated polynucleotide comprises a complement to a sequence that encodes a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 20, and conservatively modified SEQ ID NO: 20. In other alternate embodiments, the nucleotide sequence may be that of degenerate variants of above-mentioned sequences. The invention also includes  
20 operably linking one or more expression control sequences to any of the above-mentioned nucleotide sequences. The invention also includes a cell comprising any of

the above-mentioned nucleotide sequences operably linked to one or more expression control sequences.

The present invention provides a substantially pure arrestin 1 polypeptide that includes amino acid sequence that contains at least a conservatively modified identity with SEQ ID NO: 2 and binds to odorant receptors. The amino acid sequence of arrestin 1 protein can differ from SEQ ID NO: 2 by non-conservative substitutions, deletions, or insertions located at positions that do not destroy the function of the arrestin 1 polypeptide. In alternate embodiments, the polypeptide has an amino acid sequence consisting of SEQ ID NO: 2. The purified polypeptide is a polypeptide that binds specifically to an antibody that binds specifically to mosquito arrestin. In other alternate embodiments, the polypeptide comprises fragments of SEQ ID NO: 2, having at least 20 consecutive residues.

The present invention also provides a substantially pure odorant receptor 1 polypeptide that includes amino acid sequence that contains at least a conservatively modified identity with SEQ ID NO: 4 and binds to arrestin. The amino acid sequence of odorant receptor 1 polypeptide can differ from SEQ ID NO: 4 by non-conservative substitutions, deletions, or insertions located at positions that do not destroy the function of the odorant receptor 1 polypeptide. In alternate embodiments, the polypeptide has an amino acid sequence consisting of SEQ ID NO: 4. In other alternate embodiments, the polypeptide comprises fragments of SEQ ID NO: 4, having at least 20 consecutive residues.

The present invention provides a substantially pure odorant receptor 2 polypeptide that includes amino acid sequence that contains at least a conservatively modified identity with SEQ ID NO: 6 and binds to arrestin. The amino acid sequence of odorant receptor 2 polypeptide can differ from SEQ ID NO: 6 by non-conservative substitutions, deletions, or insertions located at positions that do not destroy the function of the odorant receptor 2 polypeptide. In alternate embodiments, the polypeptide has an amino acid sequence consisting of SEQ ID NO: 6. In other alternate embodiments, the polypeptide comprises fragments of SEQ ID NO: 6, having at least 20 consecutive residues.

The present invention also provides a substantially pure odorant receptor 3 polypeptide that includes amino acid sequence that contains at least a conservatively modified identity with SEQ ID NO: 8 and binds to arrestin. The amino acid sequence of odorant receptor 3 polypeptide can differ from SEQ ID NO: 8 by non-conservative substitutions, deletions, or insertions located at positions that do not destroy the function of the odorant receptor 3 polypeptide. In alternate embodiments, the polypeptide has an amino acid sequence consisting of SEQ ID NO: 8. In other alternate embodiments, the polypeptide comprises fragments of SEQ ID NO: 8, having at least 20 consecutive residues.

The present invention also provides a substantially pure odorant receptor 4 polypeptide that includes amino acid sequence that contains at least a conservatively modified identity with SEQ ID NO: 14 and binds to arrestin. The amino acid sequence

of odorant receptor 4 polypeptide can differ from SEQ ID NO: 14 by non-conservative substitutions, deletions, or insertions located at positions that do not destroy the function of the odorant receptor 4 polypeptide. In alternate embodiments, the polypeptide has an amino acid sequence consisting of SEQ ID NO: 14. In other  
5 alternate embodiments, the polypeptide comprises fragments of SEQ ID NO: 14, having at least 20 consecutive residues.

The present invention also provides a substantially pure odorant receptor 5 polypeptide that includes amino acid sequence that contains at least a conservatively modified identity with SEQ ID NO: 16 and binds to arrestin. The amino acid sequence  
10 of odorant receptor 5 polypeptide can differ from SEQ ID NO: 16 by non-conservative substitutions, deletions, or insertions located at positions that do not destroy the function of the odorant receptor 5 polypeptide. In alternate embodiments, the polypeptide has an amino acid sequence consisting of SEQ ID NO: 16. In other alternate embodiments, the polypeptide comprises fragments of SEQ ID NO: 16,  
15 having at least 20 consecutive residues.

The present invention also provides a substantially pure odorant receptor 6 polypeptide that includes amino acid sequence that contains at least a conservatively modified identity with SEQ ID NO: 18 and binds to arrestin. The amino acid sequence  
20 of odorant receptor 6 polypeptide can differ from SEQ ID NO: 18 by non-conservative substitutions, deletions, or insertions located at positions that do not destroy the function of the odorant receptor 6 polypeptide. In alternate embodiments, the



polypeptide has an amino acid sequence consisting of SEQ ID NO: 18. In other alternate embodiments, the polypeptide comprises fragments of SEQ ID NO: 18, having at least 20 consecutive residues.

The present invention also provides a substantially pure odorant receptor 7 polypeptide that includes amino acid sequence that contains at least a conservatively modified identity with SEQ ID NO: 20 and binds to arrestin. The amino acid sequence of odorant receptor 7 polypeptide can differ from SEQ ID NO: 20 by non-conservative substitutions, deletions, or insertions located at positions that do not destroy the function of the odorant receptor 7 polypeptide. In alternate embodiments, the polypeptide has an amino acid sequence consisting of SEQ ID NO: 20. In other alternate embodiments, the polypeptide comprises fragments of SEQ ID NO: 20, having at least 20 consecutive residues.

The invention also provides an arrestin 1 antibody, which comprises polyclonal or monoclonal antibodies. The antibody can be conjugated to a detectable label.

Another aspect of the present invention provides an odorant receptor 1 antibody, which comprises polyclonal or monoclonal antibodies. The antibody can be conjugated to a detectable label. Antibody labels and methods are well known in the art.

The present invention also provides an odorant receptor 2 antibody, which comprises polyclonal or monoclonal antibodies. The antibody can be conjugated to a detectable label.

Another aspect of the present invention provides an odorant receptor 3  
antibody, which comprises polyclonal or monoclonal antibodies. The antibody can be  
conjugated to a detectable label.

Another aspect of the present invention provides an odorant receptor 4  
5 antibody, which comprises polyclonal or monoclonal antibodies. The antibody can be  
conjugated to a detectable label.

Another aspect of the present invention provides an odorant receptor 5  
antibody, which comprises polyclonal or monoclonal antibodies. The antibody can be  
conjugated to a detectable label.

Another aspect of the present invention provides an odorant receptor 6  
10 antibody, which comprises polyclonal or monoclonal antibodies. The antibody can be  
conjugated to a detectable label.

Another aspect of the present invention provides an odorant receptor 7  
antibody, which comprises polyclonal or monoclonal antibodies. The antibody can be  
15 conjugated to a detectable label.

The present invention also presents a method of producing arrestin 1 protein.  
The method includes the following steps: (a) providing a cell transformed with an  
isolated DNA comprising a nucleotide sequence that encodes an amino acid sequence  
of SEQ ID NO: 2; (b) culturing the cell; and (c) collecting from the cell or the medium  
20 of the cell the polypeptide encoded by the polynucleotide sequence. Certain

alternatives to SEQ ID NO: 2 are described above (*e.g.* conservative variants and hybridization variants).

The present invention also provides a method of manufacturing odorant receptor 1 protein. The method includes the following steps: (a) providing a cell transformed with an isolated DNA comprising a nucleotide sequence that encodes an amino acid sequence of SEQ ID NO: 4; (b) culturing the cell; and (c) collecting from the cell or the medium of the cell the polypeptide encoded by the polynucleotide sequence.

The present invention provides a method of manufacturing odorant receptor 2 protein. The method includes the following steps: (a) providing a cell transformed with an isolated DNA comprising a nucleotide sequence that encodes an amino acid sequence of SEQ ID NO: 6; (b) culturing the cell; and (c) collecting from the cell or the medium of the cell the polypeptide encoded by the polynucleotide sequence.

The present invention also provides a method of manufacturing odorant receptor 3 protein. The method includes the following steps: (a) providing a cell transformed with an isolated DNA comprising a nucleotide sequence that encodes an amino acid sequence of SEQ ID NO: 8; (b) culturing the cell; and (c) collecting from the cell or the medium of the cell the polypeptide encoded by the polynucleotide sequence.

The present invention also provides a method of manufacturing odorant receptor 4 protein. The method includes the following steps: (a) providing a cell transformed with an isolated DNA comprising a nucleotide sequence that encodes an amino acid sequence of SEQ ID NO: 14; (b) culturing the cell; and (c) collecting from

the cell or the medium of the cell the polypeptide encoded by the polynucleotide sequence.

The present invention also provides a method of manufacturing odorant receptor 5 protein. The method includes the following steps: (a) providing a cell transformed with an isolated DNA comprising a nucleotide sequence that encodes an amino acid sequence of SEQ ID NO: 16; (b) culturing the cell; and (c) collecting from the cell or the medium of the cell the polypeptide encoded by the polynucleotide sequence.

The present invention also provides a method of manufacturing odorant receptor 6 protein. The method includes the following steps: (a) providing a cell transformed with an isolated DNA comprising a nucleotide sequence that encodes an amino acid sequence of SEQ ID NO: 18; (b) culturing the cell; and (c) collecting from the cell or the medium of the cell the polypeptide encoded by the polynucleotide sequence.

The present invention also provides a method of manufacturing odorant receptor 7 protein. The method includes the following steps: (a) providing a cell transformed with an isolated DNA comprising a nucleotide sequence that encodes an amino acid sequence of SEQ ID NO: 20; (b) culturing the cell; and (c) collecting from the cell or the medium of the cell the polypeptide encoded by the polynucleotide sequence.

The present invention also provides a method for identifying a mosquito olfaction molecule binding compound. The method includes the following steps: (a) providing an isolated mosquito olfaction molecule; (b) contacting a test agent with the isolated mosquito olfaction molecule; and (c) detecting whether the test agent is bound to the isolated mosquito olfaction molecule. Methods of detection are well known in the art. In certain embodiments, the isolated mosquito olfaction molecule further comprises a polypeptide having an amino acid sequence as set forth in SEQ ID NO: 2 or variants thereof as described herein (As used herein this statement means conservatively modified variants, hybridization variants, and variants to which antibodies bind specifically). In alternate embodiments, the isolated mosquito olfaction molecule further comprises a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO. 4, SEQ ID NO. 6, SEQ ID NO. 8, SEQ ID NO. 14, SEQ ID NO. 16, SEQ ID NO. 18, SEQ ID NO. 20, conservatively modified SEQ ID NO: 4, conservatively modified SEQ ID NO: 6, conservatively modified SEQ ID NO: 8, conservatively modified SEQ ID NO: 14, conservatively modified SEQ ID NO: 16, conservatively modified SEQ ID NO: 18, and conservatively modified SEQ ID NO: 20. In other embodiments, contacting the test agent with the isolated mosquito olfaction molecule further comprises contacting under native conditions. In alternate embodiments, detecting specific binding of the test agent to the isolated mosquito olfaction molecule further comprises immunoprecipitation.

The present invention also presents a screening method for identifying a compound that inhibits binding of mosquito arrestin to a mosquito odorant receptor. The method includes the following steps: (a) providing an antibody that binds to an isolated mosquito olfaction molecule; (b) providing a mosquito olfaction molecule binding compound; (c) providing a test sample comprising the mosquito arrestin polypeptide and mosquito odorant receptor; (d) combining the mosquito olfaction molecule binding compound, the antibody, and the test sample in reaction conditions that allow a complex to form in the absence of the mosquito olfaction molecule binding compound., wherein the complex includes the antibody, mosquito arrestin and mosquito odorant receptor; and (e) determining whether the mosquito olfaction molecule binding compound decreases the formation of the complex, wherein a decrease indicates that the mosquito olfaction molecule binding compound is a compound that inhibits the binding of mosquito arrestin to mosquito odorant receptor. In certain embodiments, the mosquito odorant receptor further comprises a polypeptide having any of the following sequences: SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, conservatively modified SEQ ID NO: 4, conservatively modified SEQ ID NO: 6, conservatively modified SEQ ID NO: 8, conservatively modified SEQ ID NO: 16, conservatively modified SEQ ID NO: 18, conservatively modified SEQ ID NO: 20 or conservatively modified SEQ ID NO: 14.

Various features and advantages of the invention will be apparent from the following detailed description and from the claims.

### BRIEF DESCRIPTION OF THE DRAWINGS

5 FIG. 1 is the nucleotide sequence (SEQ ID NO: 1) of arrestin 1 isolated from *Anopheles gambiae*.

FIG. 2 is the deduced amino acid sequence of arrestin 1 isolated from *Anopheles gambiae* (SEQ ID NO: 2).

10 FIG. 3a-b are the nucleotide sequence (SEQ ID NO: 9) and deduced amino acid sequence (SEQ ID NO: 4) of odorant receptor 1 isolated from *Anopheles gambiae*.

FIG. 4a-b are the nucleotide sequence (SEQ ID NO: 10) and deduced amino acid sequence (SEQ ID NO: 6) of odorant receptor 2 isolated from *Anopheles gambiae*.

FIG. 5a-b are the nucleotide sequence (SEQ ID NO: 11) and deduced amino acid sequence (SEQ ID NO: 8) of odorant receptor 3 isolated from *Anopheles gambiae*.

15 FIG. 6a-b are the nucleotide sequence (SEQ ID NO: 12) and deduced amino acid sequence (SEQ ID NO: 14) of odorant receptor 4 isolated from *Anopheles gambiae*.

FIG. 7 is a table of preferred codons used to deduce amino acid sequences from nucleotide sequences for *Anopheles gambiae*.

20 FIG. 8 is a table listing cDNA and polypeptide sequences with corresponding SEQ ID numbers and Figure numbers.

FIG. 9a-b are the nucleotide sequence (SEQ ID NO: 21) and deduced amino acid sequence (SEQ ID NO: 16) of odorant receptor 5 isolated from *Anopheles gambiae*.

FIG. 10a-b are the nucleotide sequence (SEQ ID NO: 22) and deduced amino acid sequence (SEQ ID NO: 18) of odorant receptor 6 isolated from *Anopheles gambiae*.

5 FIG. 11a-b are the nucleotide sequence (SEQ ID NO: 23) and deduced amino acid sequence (SEQ ID NO: 20) of odorant receptor 7 isolated from *Anopheles gambiae*.

## DETAILED DESCRIPTION OF THE INVENTION

Arrestins interact with odorant receptors to cause changes in cellular function. Interruption of normal arrestin function will lead to over stimulation of the olfaction system. Consequently, substances that block the arrestin - odorant receptor interaction can interfere with a mosquito's ability to home in on sources of bloodmeal, such as humans. Screening for substances that modulate arrestin - odorant receptor interaction is therefore useful for identifying pest control agents and for treatment of malaria. The deduced amino acid sequence and arrestin contains several domains implicated in arrestin function. The motifs potentiation consensus Src homology 3 (SH3) binding sites. Cohen, *et al.*, 1995, Cell, 80:237. Sequence comparisons with the DDBJ/EMBL/GenBank and SWISSPROT databases were performed using the GCG software. Devereux, *et al.*, 1984, Nucleic Acids Res., 12:387-395. Protein alignment was also performed using the Clustal W software package. Thompson, *et al.*, 1994, Nucleic Acids Res, 22:4673-4680. Additionally,



arrestin has been submitted to the GenBank database with accession No.  
AY017417.

As used herein, "native conditions" means natural conditions as found within  
the ordinary conditions found within *Anopheles gambiae*.

5 As used herein, "stringent conditions" means the following: hybridization at  
42° C in the presence of 50% formamide; a first wash at 65° C with about 2 x SSC  
containing 1% SDS; followed by a second wash at 65° C with 0.1 x SSC. Salt  
concentrations and temperature may be modified. Such modifications may be found  
in Sambrook *et al.*, 1989, Molecular Cloning: A Laboratory Manual (2nd Edition),  
10 Cold Spring Harbor Press, Cold Spring Harbor, N.Y. The hybridizing part of the  
nucleic acid is generally at least 15 nucleotides in length.

As used herein, "purified polypeptide" means a polypeptide that is  
substantially free from compounds normally associated with the polypeptide in the  
natural state. The absence of such compounds may be determined by detection of  
15 protein bands subsequent to SDS-PAGE. Purity may also be assessed in other ways  
known to those of ordinary skill in the art. The term, as defined herein, is not  
intended to exclude (1) synthetic or artificial combinations of the polypeptides with  
other compounds, (2) polypeptides having minor impurities which do not interfere  
with biological activity.

20 As used herein, "isolated polynucleotide" means a polynucleotide having a  
structure that is not identical to any naturally occurring nucleic acid or of any

fragment of a naturally occurring genomic nucleic acid spanning more than three separate genes. Thus, the term includes (1) a nucleic acid incorporated into a vector or into the genomic DNA of a prokaryote or eukaryote in a manner such that the resulting molecule is not identical to any naturally occurring vector or genomic DNA; (2) a separate molecule of a cDNA, a genomic fragment, a fragment produced by polymerase chain reaction (PCR), or a restriction fragment; and (3) a recombinant nucleotide sequence that is part of a gene encoding a fusion protein. This definition of "isolated polynucleotide" supersedes and controls all other definitions known in the art.

As used herein, "hybridization probe" means nucleic acid that is labeled for detection, such as labeling with radiation. Hybridization probes are well known in the art.

As used herein, "culturing the cell" means providing culture conditions that are conducive to polypeptide expression. Such culturing conditions are well known in the art.

As used herein, "operably linked" means incorporated into a genetic construct so that expression control sequences effectively control expression of a gene of interest.

As used herein, "protein" means any peptide-linked chain of amino acids, regardless of length or post-translational modification, *e.g.*, glycosylation or phosphorylation.

As used herein, "sequence identity" means the percentage of identical subunits at corresponding positions in two sequences when the two sequences are aligned to maximize subunit matching, i.e., taking into account gaps and insertions. When a subunit position in both of the two sequences is occupied by the same monomeric subunit, *e.g.*, if a given position is occupied by an adenine in each of two DNA molecules, then the molecules are identical at that position. For example, if 7 positions in a sequence 10 nucleotides in length are identical to the corresponding positions in a second 10-nucleotide sequence, then the two sequences have 70% sequence identity. Preferably, the length of the compared sequences is at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 100 nucleotides. Sequence identity is typically measured using sequence analysis software (*e.g.*, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705).

As used herein, "mosquito olfaction molecule" means a polypeptide that is involved in the modulation of the mosquito olfaction system. By way of illustration, and not limitation, mosquito olfaction molecules have the following characteristics: (1) G protein-coupled seven-transmembrane domain receptors, (2) sequence conservation regarding positions of a subset of introns and the length of the deduced protein, (3) they are selectively expressed in olfactory receptor neurons, and (4) they have highly conserved structural motifs. Odorant receptors 3, 4 and 5 are clustered



olfaction molecules additionally include polypeptides having the characteristics noted in the definition of the term.

As used herein, "mosquito olfaction molecule-specific antibody" means an antibody that binds to a mosquito olfaction molecule. The term includes polyclonal and monoclonal antibodies.

As used herein, "substantially pure protein" means a protein separated from components that naturally accompany it. Typically, the protein is substantially pure when it is at least 60%, by weight, free from the proteins and other naturally-occurring organic molecules with which it is naturally associated. In certain embodiments, the purity of the preparation is at least 75%, more preferably at least 90%, 95% and most preferably at least 99%, by weight. A substantially pure mosquito olfaction molecule protein can be obtained, for example, by extraction from a natural source, by expression of a recombinant nucleic acid encoding a mosquito olfaction molecule polypeptide, or by chemical synthesis. Purity can be measured by any appropriate method, e.g., column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis. A chemically-synthesized protein or a recombinant protein produced in a cell type other than the cell type in which it naturally occurs is, by definition, substantially free from components that naturally accompany it. Accordingly, substantially pure proteins include those having sequences derived from eukaryotic organisms but synthesized in *E. coli* or other prokaryotes.

As used herein, "fragment", as applied to a polypeptide (e.g., arrestin 1 polypeptide), means at least about 10 amino acids, usually about 20 contiguous amino acids, preferably at least 40 contiguous amino acids, more preferably at least 50 amino acids, and most preferably at least about 60 to 80 or more contiguous amino acids in length. Such peptides can be generated by methods known to those skilled in the art, including proteolytic cleavage of the protein, de novo synthesis of the fragment, or genetic engineering.

As used herein, "test sample" means a sample that contains arrestin 1, or conservatively modified variant thereof, in combination with at least one of the following: odorant receptor 1, odorant receptor 2, odorant receptor 3, odorant receptor 5, odorant receptor 6, odorant receptor 7, odorant receptor 4, conservatively modified variants of the above, or other odorant receptors known in the art.

As used herein, "vector" means a replicable nucleic acid construct, e.g., a plasmid or viral nucleic acid. Preferably, expression is controlled by an expression control sequence.

As used herein, "conservatively modified" applies to both amino acid and nucleic acid sequences. Regarding nucleic acid sequences, conservatively modified refers to those nucleic acids which encode identical or conservatively modified variants of the amino acid sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For example, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine.

Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of ordinary skill  
5 will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine; and UGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide of the present invention is implicit in each described polypeptide sequence and incorporated  
10 herein by reference.

As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified  
15 variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Thus, any number of amino acid residues selected from the group of integers consisting of from 1 to 15 can be so altered. Thus, for example, 1, 2, 3, 4, 5, 7, or 10 alterations can be made. Conservatively modified variants typically provide similar biological activity as the unmodified polypeptide  
20 sequence from which they are derived. For example, substrate specificity, enzyme activity, or ligand/receptor binding is generally at least 30%, 40%, 50%, 60%, 70%,

80%, or 90% of the native protein for its native substrate. Conservative substitution tables providing functionally similar amino acids are well known in the art. The following six groups each contain amino acids that are conservative substitutions for one another: 1) Alanine (A), Serine (S), Threonine (T); 2) Aspartic acid (D),  
5 Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W). See also, Creighton (1984) Proteins W.H. Freeman and Company.

As used herein, "immunogenic fragment" means the fragment of a polypeptide that is capable of eliciting an immunogenic response.  
10

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the  
15 present invention, the preferred methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present document, including definitions, will control. Unless otherwise indicated, materials, methods, and examples described herein are illustrative only and not intended to be  
20 limiting.



## Structure and Function

The genes disclosed herein have homology to corresponding arrestin and odorant receptor *Drosophila melanogaster* genes. Fox, *et al.*, 2001, PNAS 98:14693-14697. The genes disclosed herein have the utility disclosed within this patent application.

A full-length *Anopheles gambiae* arrestin 1 cDNA has been cloned and sequenced. The arrestin 1 cDNA clone contains 1964 bp and includes a complete open reading frame that encodes a protein 383 amino acids in length, as seen in Figure 1. The open reading frame from the methionine includes 383 amino acids, yielding a slightly basic polypeptide (PI=8.0) with a predicted molecular weight of 42.8 KD.

A full-length *Anopheles gambiae* odorant receptor 1 genomic DNA has been sequenced. The odorant receptor 1 genomic DNA contains 3895 bp and includes a deduced open reading frame that encodes a protein 394 amino acids in length.

A full-length *Anopheles gambiae* odorant receptor 2 genomic DNA has been sequenced. The odorant receptor 2 genomic DNA contains 4985 bp and includes a deduced open reading frame that encodes a protein 380 amino acids in length.

A full-length *Anopheles gambiae* odorant receptor 3 genomic DNA has been sequenced. The odorant receptor 3 genomic DNA contains 2083 bp and includes a deduced open reading frame that encodes a protein 411 amino acids in length.

A full-length *Anopheles gambiae* odorant receptor 4 genomic DNA has been sequenced. The odorant receptor 4 genomic DNA contains 2374 bp and includes a deduced open reading frame that encodes a protein 394 amino acids in length.

A full-length *Anopheles gambiae* odorant receptor 5 genomic DNA has been  
5 sequenced. The odorant receptor 5 genomic DNA contains 2272 bp and includes a deduced open reading frame that encodes a protein 391 amino acids in length.

A partial *Anopheles gambiae* odorant receptor 6 genomic DNA has been sequenced. The odorant receptor 6 genomic DNA contains 931 bp and includes a deduced open reading frame that encodes a protein 157 amino acids in length.

A full-length *Anopheles gambiae* odorant receptor 7 genomic DNA has been  
10 sequenced. The odorant receptor 7 genomic DNA contains 11,103 bp and includes a deduced open reading frame that encodes a protein 401 amino acids in length.

#### Expression Control Sequences and Vectors

The mosquito olfaction molecules of this invention can be used in a method to identify a mosquito olfaction molecule binding compound. If desired, the mosquito olfaction molecule binding compounds may be further tested for ability to inhibit binding of arrestin to an odorant receptor. Methods for this test are described  
15  
20 herein. In certain embodiments, the DNA that encodes the arrestin 1 polypeptide ("ARR1 DNA") may be cloned into an expression vector, i.e., a vector wherein ARR1

10056405-012402  
20420-9499007

DNA is operably linked to expression control sequences. The need for expression control sequences will vary according to the type of cell in which the ARR1 DNA is to be expressed. Generally, expression control sequences include a transcriptional promoter, enhancer, suitable mRNA ribosomal binding sites, and sequences that terminate transcription and translation. One of ordinary skill in the art can select proper expression control sequences. Standard methods can be used by one skilled in the art to construct expression vectors. See generally, Sambrook *et al.*, 1989, Molecular Cloning: A Laboratory Manual (2nd Edition), Cold Spring Harbor Press, Cold Spring Harbor, N.Y. Vectors useful in this invention include, but are not limited to plasmid vectors and viral vectors.

All other nucleic acid sequences disclosed herein may also be operably linked to expression control sequences. The expression control sequences described above may be used. As mentioned above, methods known to those of ordinary skill in the art may be used to insert nucleic acid sequences into expression control sequences.

Methods known to those of ordinary skill in the art may be used to introduce the nucleic acid and expression control sequence into eukaryotic and/or prokaryotic cells. An example of prokaryotic cells is BL21 (DE3)pLysS bacteria. An example of eukaryotic cells is Sf9.

In certain embodiments of the invention, ARR1 DNA is introduced into, and expressed in, a prokaryotic cell, *e.g.*, BL21 (DE3)pLysS bacteria.

In certain embodiments of the invention, the ARR1 DNA is introduced into, and expressed in, a eukaryotic cell *in vitro*. Eukaryotic cells useful for expressing ARR1 DNA *in vitro* include, but are not limited to Sf9 cells. Transfection of the eukaryotic cell can be transient or stable.

5

#### Mosquito Olfaction Molecule-Specific Antibody

An animal is immunized with a mosquito olfaction molecule (e.g., arrestin 1 polypeptide). The animal produces antibodies to the mosquito olfaction molecule. The production and collection of the polyclonal antibodies was performed by Lampire Biological Laboratories, Inc. of Pipersville, PA 18947, using techniques known in the art.

10

#### Mosquito Olfaction Molecule Antibody Label

15

In some embodiments of the invention, the mosquito olfaction molecule-specific antibody includes a detectable label. Many detectable labels can be linked to, or incorporated into, an antibody of this invention. The following are examples of useful labels: radioactive, non-radioactive isotopic, fluorescent, chemiluminescent, paramagnetic, enzyme, or colorimetric.

20

Examples of useful enzyme labels include malate hydrogenase, staphylococcal dehydrogenase, delta-5-steroid isomerase, alcohol dehydrogenase, alpha-glycerol phosphate dehydrogenase, triose phosphate isomerase, peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, and glucoamylase, acetylcholinesterase. Examples of useful radioisotopic labels include  $^3\text{H}$ ,  $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ , and  $^{14}\text{C}$ . Examples of useful fluorescent labels include fluorescein, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, and fluorescamine. Examples of useful chemiluminescent label types include luminal, isoluminal, aromatic acridinium ester, imidazole, acridinium salt, oxalate ester, luciferin, luciferase, and aequorin.

Antibody labels can be coupled to, or incorporated into antibodies by use of common techniques known to those of ordinary skill in the art. Typical techniques are described by Kennedy *et al.*, 1976, Clin. Chim. Acta, 70:1-31; and Schurs *et al.*, 1977, Clin. Chim. Acta, 81: 1-40. Useful chemical coupling methods include those that use glutaraldehyde, periodate, dimaleimide and m-maleimido-benzyl-N-hydroxy-succinimide ester.

Screening assays

The present invention provides, in part, a screen for mosquito olfaction molecule binding compounds with the ability to interrupt the interaction of arrestin with an odorant receptor. Identifying that a test agent will bind a mosquito olfaction molecule is one part. Once a test agent has demonstrated its ability to bind a mosquito olfaction molecule, it is properly called a mosquito olfaction molecule binding compound. Since it is possible for a mosquito olfaction molecule binding compound to bind without necessarily interrupting the arrestin-odorant receptor interaction, it is proper to further assay in order to determine that the interaction is disrupted. The ability of the mosquito olfaction molecule binding compound to interrupt the arrestin-odorant receptor interaction may be assayed.

In certain embodiments, a test agent is identified as a mosquito olfaction molecule binding compound by the following method. One of the mosquito olfaction molecules is immobilized (*e.g.*, arrestin 1). Polypeptides can be immobilized using methods known in the art. Such methods include the use of Affigel (Biorad) or activated agarose or sepharose to which significant amounts of polypeptides can be directly coupled. The immobilized polypeptide (*e.g.*, arrestin 1) is contacted with the test agent. Unbound test agent can be removed by washing with binding buffer. Then, the bound test agent is eluted by a salt gradient. The material that is bound to the immobilized polypeptide may be purified by SDS-PAGE. Other methods known by one of ordinary skill in the art for identifying an interaction between two

proteins include affinity purification, co-immunoprecipitation, and far-western blotting.

In certain embodiments, the following method is used to screen for substances capable of interrupting arrestin-odorant receptor interaction. The following method of detecting protein-protein interaction will also provide information regarding the lack of protein-protein interactions. The two-hybrid method is a well known genetic assay used to detect protein-protein interactions *in vivo*. See, *e.g.*, Bartel *et al.*, 1993, In Cellular Interactions in Development: A Practical Approach, Oxford University Press, Oxford, pp. 153-179; Chien *et al.*, 1991, Proc. Natl. Acad. Sci. USA, 88:9578-9582; Fields *et al.*, 1989, Nature, 340:245-247; Fritz *et al.*, 1992, Curr. Biol., 2:403-405; Guarente, L., 1993, Proc. Natl. Acad. Sci. USA, 90:1639-1641. There are multiple combinations available between arrestin and the seven odorant receptors. A GAL4 binding domain is linked to an arrestin fragment (*e.g.*, arrestin 1 polypeptide) and a GAL4 transactivation domain is linked to an odorant receptor fragment (*e.g.*, odorant receptor 1 polypeptide). A GAL4 binding site is linked to a reporter gene such as lacZ. All three elements are contacted in the presence and absence of a mosquito olfaction molecule binding compound. The level of expression of the reporter gene is monitored. A decrease in the level of expression of lacZ means that the mosquito olfaction molecule binding compound interrupts the interaction of arrestin with the odorant receptor.

In an alternate embodiment, the following is a method that will identify whether a mosquito olfaction molecule binding compound will interrupt the interaction between arrestin and an odorant receptor. The following method of co-immunoprecipitation may make use of the available panel of antibodies to any arrestin or odorant receptor. Since this method makes use of antibodies that demonstrate the ability to immunoprecipitate the mosquito olfaction molecule and other proteins to which it is bound, the ability of a mosquito olfaction molecule binding compound to inhibit the interaction of the mosquito olfaction molecule will serve as the measure of the compound's interruption ability.

Also disclosed herein is a method of modulating arrestin 1 biological activity. In certain embodiments, the method comprises administering an arrestin 1 biological activity-modulating amount of a mosquito olfaction molecule binding compound. Upon administration, arrestin 1 is contacted with the mosquito olfaction molecule binding compound. Such contact results in modulating arrestin 1 biological activity. The mosquito olfaction molecule binding compound may be administered as an aerosol, solid, or liquid, such that delivery occurs through contact with the body of the target subject. For example, administration may occur by absorption through the exterior surfaces of the target subject, ie. mosquitoes, or by intake through other apertures of the target subject [proboscis (or other feeding aperture), or spiracles (or other respiratory apertures)]. An activity-modulating amount of mosquito olfaction molecule binding compound is an amount that is



sufficient to prohibit at least about 50% of the arrestin 1 (SEQ ID NO: 2) molecules from interacting with any odorant receptors.

All citations and references described in this patent application are hereby incorporated herein by reference, in their entirety. Also incorporated in this specification are the exhibits filed herewith. The present invention is further illustrated by the following specific examples. The examples are provided for illustration only and are not to be construed as limiting the scope or content of the invention in any way.

### Example 1

#### Protein expression

A cDNA encoding arrestin 1 is subcloned into the pBlueScript II (KS) vector (Novagen, Madison, WI) at the BamHI/NdeI restriction sites for DNA sequencing. The cDNA encoding arrestin 1 is subsequently subcloned into the bacterial expression plasmid pET15b (Novagen, Madison, WI). The bacterial expression plasmid containing the arrestin 1 cDNA is transformed into BL21 (DE3)pLysS bacteria (Novagen, Madison, WI) for high levels of arrestin 1 expression. Methods are known in the art for isolating the expressed protein.

Expression of other nucleic acids disclosed herein is achieved by using the above-referenced method. Once the odorant receptor is in protein form, it may be used as described within this application.

## Example 2

### Mosquito Olfaction Molecule Specific Antibody

The cDNA encoding arrestin 1 is subcloned into the bacterial expression plasmid pET15b (Novagen, Madison, WI). The vector is transformed into BL21 (DE3)pLysS bacteria (Novagen, Madison, WI) for high levels of arrestin 1 expression. Rapid purification is performed using His-Bind affinity Resin (Novagen, Madison, WI). Native recombinant arrestin 1 is then denatured using gel purification on SDS-polyacrylamide gel electrophoresis followed by staining with 0.05% Coomassie Brilliant Blue (Sigma-Aldrich, St. Louis, MO). Polyclonal antibodies were generated in rabbits by Lampire Biological Laboratories, Inc. of Pipersville, PA 18947. Polyclonal antibodies may be generated for any of the odorant receptors disclosed herein.

## Example 3

### Identification of a mosquito olfaction molecule binding compound

Arrestin 1 polypeptide is expressed in and purified from BL21 (DE3)pLysS bacteria (Novagen, Madison, WI). Arrestin 1 is incubated with a test agent in Phosphate Buffered Saline (pH 7.5), 0.1% Tween-20, and 0.1% broad spectrum protease inhibitors for 90 minutes at 4° C. Anti-arrestin 1 polyclonal sera is added to the reaction at a dilution of 1:2000 and incubated for an additional 60 minutes. The complexes, consisting of either polypeptide-antibody or test agent-polypeptide-antibody are isolated by the addition of  $1 \times 10^7$  Dynalbeads M280 (sheep anti-Rabbit

2042T0-649901  
IgG) followed by incubation at the same temperature for an additional 60 minutes.  
Isolation of the complexes is completed by using the DYNAL Magnetic Particle  
Concentrator (DynaL Inc., Lake Success, NY). The complexes are washed three  
times with broad spectrum protease inhibitors. Content of the complexes is assayed  
5 by SDS-PAGE followed by silver staining and western blotting. Common methods  
are known by those of ordinary skill in the art for silver staining and western  
blotting. See generally, Sambrook *et al.*, 2001, Molecular Cloning: A Laboratory  
Manual (3rd Edition), Cold Spring Harbor Press, Cold Spring Harbor, N.Y.  
Obviously, the presence of the test agent, polypeptide, and antibody indicates that  
10 the test agent binds to the polypeptide.

#### Example 4

##### Identification of a compound that inhibits binding of arrestin to an odorant receptor

15 Arrestin 1 polypeptide and odorant receptor 1 polypeptide are expressed in  
and purified from BL21 (DE3)pLysS bacteria (Novagen, Madison, WI). Arrestin 1  
polypeptide and odorant receptor 1 polypeptide are incubated with a mosquito  
olfaction molecule binding compound in Phosphate Buffered Saline (pH 7.5), 0.1%  
Tween-20, and 0.1% broad spectrum protease inhibitors for 90 minutes at 4° C.  
20 Anti-arrestin 1 polyclonal sera is added to the reaction at a dilution of 1:2000 and  
incubated for an additional 60 minutes. The complexes, consisting of either

antibody-arrestin 1-odorant receptor 1 or antibody-arrestin 1, are isolated by the addition of  $1 \times 10^7$  Dynalbeads M280 (sheep anti-Rabbit IgG) followed by incubation at the same temperature for an additional 60 minutes (Dynal Inc., Lake Success, NY). Once the isolation of the complexes is completed by using the DYNAL  
5 Magnetic Particle Concentrator, (Dynal Inc., Lake Success, NY), the complexes are washed three times with broad spectrum protease inhibitors. The content of the complexes is assayed by SDS-PAGE followed by silver staining and western blotting. Common methods are known by those of ordinary skill in the art for silver staining and western blotting. See generally, Sambrook *et al.*, 2001, Molecular  
10 Cloning: A Laboratory Manual (3rd Edition), Cold Spring Harbor Press, Cold Spring Harbor, N.Y.

### Example 5

#### Far western blotting to analyze components of a protein mixture

15 The protein sample is fractionated on an SDS-PAGE gel. After electrophoresis at a voltage and time that is known in the art, the proteins are transferred from the gels onto a solid support membrane by electroblotting. Transferred membranes may be stained with Ponceau S to facilitate location and identification of specific proteins. Nonspecific sites on the membranes are blocked  
20 with standard blocking reagents, and the membranes are then incubated with a

radiolabeled non-antibody protein probe. After washing, proteins that bind to the probe are detected by autoradiography.

The content of the solutions used within this protocol are disclosed in Wiley's Current Protocols in Cell Biology.

5 The protein sample to be analyzed is resuspended in 1x SDS sample buffer. Approximately 50 to 100 ug can be loaded in each lane of the gel. The samples are separated with SDS-PAGE. The proteins are transferred to nitrocellulose by electroblotting.

10 After transfer, stain the membrane for 5 min in ~100 ml freshly diluted 1x Ponceau S staining solution. The membrane is then destained by washing it in several changes of deionized water until the proteins are clearly visible. Continue to destain for an additional 5 min in water until the red staining fades.

15 The membrane is then blocked for 2 hr in 200 ml blocking buffer I at room temperature with gentle agitation. Incubate the membrane in 200 ml of blocking buffer II for 2 hours and rinse the membrane briefly in 100 ml of 1 x PBS.

Prior to probing, the membrane is preincubated for 10 min in 50 ml of 1x probe dilution buffer without the probe at room temperature. The probe is added to the membrane and incubated for 2 hours at room temperature. The membrane is washed with 200 ml 1x PBS for 5 min, room temperature. Repeat the wash step  
20 three additional times. Air dry the filter and expose to x-ray film with intensifying screen. An overnight exposure is typically sufficient.

### Sequence ID Listing

#### SEQ ID NO:1

cDNA Nucleic Acid Sequence

1964 nucleotides

5 Mosquito arrestin 1

ACAGGAACGACGGTTGTGATCCCTCCACTGGTGGTGACACGAATCATAAGCAT  
TATTTTCATACCTAAAAAACAAAATCTACAAAAAAAAGCTTCATTCCCATCGAAA  
AAACTTTCTTGTGAAATCAACCGAGCTAACAAACAACATCCTGTGCAAAATCTA  
10 GCAGTGAAAGTGTGATATCGTATACCTGTACCTGTAAACCGTTGTGCGCGTGT  
GTGCCTTTTGTGTATCAATTTTGTGGAAAACAGAAAATACATCAAAATGGTTTAC  
AATTTCAAAGTCTTCAAGAAGTGCGCCCCTAATGGAAAGGTTACGCTGTACATG  
GGCAAGCGTGACTTTGTAGACCACGTTTCCGGCGGTTGAACCGATCGATGGTAT  
CGTCGTCCTCGATGATGAGTACATTCGTGACAACCGTAAGGTATTCGGTCAGAT  
15 TGTCTGCAGTTTCCGCTACGGCCGCGAAGAGGACGAGGTGATGGGACTAAACT  
TCCAGAAGGAGTTATGCCTCGCTTCCGAACAGATCTACCCGCGTCCGGAAAAG  
TCGGACAAGGAGCAGACCAAGCTCCAGGAGCGACTGCTGAAGAAGCTGGGTTC  
GAACGCCATCCCGTTCACGTTCAACATCTCGCCGAATGCTCCGTCTTCGGTCAC  
GCTGCAGCAGGGCGAAGATGATAATGGAGACCCGTGCGGTGTGTCTGCTACTACG  
20 TGAAGATCTTTGCCGGTGAGTCGGAAACCGATCGTACGCACCGTCGCAGCACC  
GTTACGCTCGGCATACGCAAGATCCAGTTCGCACCGACCAAGCAGGGCCAGCA  
GCCGTGCACGCTGGTGCGCAAGGACTTTATGCTAAGCCCGGGAGAGCTGGAGC  
TCGAGGTCACACTAGACAAGCAGCTGTACCTGCACGGGGAGCGAATAGGCGTC  
AACATCTGCATCCGCAACAACCTCGAACAAAATGGTCAAGAAGATTAAGGCCAT  
25 GGTCCAGCAGGGTGTGGATGTGGTGCTGTTCCAGAATGGTAGCTACCGCAACA  
CAGTGGCATCGCTGGAGACTAGCGAGGGTTGCCCAATTCAGCCCGGCTCCAGT  
CTGCAGAAGGTAATGTACCTCACGCCGCTGCTGTCCTCGAACAAGCAGCGACG  
TGGCATCGCCCTGGACGGTCAGATCAAGCGTCAGGATCAGTGTTTGGCCTCGA  
CAACCCTCTTGGCTCAACCGGATCAGCGAGATGCTTTTCGGCGTTATCATATCGT  
30 ATGCCGTAAAGGTTAAGCTTTTCCCTCGGCGCACTCGGCGGCGAGCTGTCGGCG  
GAACTTCCATTTGTGCTGATGCACCCAAAGCCCGGCACCAAGGCTAAGGTCAT  
CCATGCCGACAGCCAGGCCGACGTAGAAACTTTCCGACAGGATACAATCGACC  
AGCAGGCATCAGTTGACTTTGAATAGACGACGCAACGGTTTGGAAATGCTACC  
TACTACCCAGGCATGGGCTAACACGACGAACGAACACTACTACTAAGCATA  
35 AAAACAGGAAAAAAAAATGGAAAACCTTAAAAAATGGATCATACAACCGAACGC  
AAACGACCTACGACGATCGATCTCACTTCCCCGTCTTTTTTCATCCTAAGCAATA  
GAACGATGGTAGAAAAGGAAGATAAAGATGGAGAGAAAGTCACGTGTATCAAT  
GACGACGACTACCAAAACTGAAGACGTAACACATGTTCCCCAGCGAGCGGTAA  
CTGTTCTGTTCTGACACCTTCCGCTCGACAATGTACCTTTTAAAAACATACAAA

TTAGAAGTCGTCTTCACTACCTTCAACCAATCCAGCCACTTTGGTATATACTTTT  
CATAGAATCCTTCTGAGCGCAAGGACCCTATTGAAATTCAGTGTTATTTTGTA  
CTGCGACCAAATGCCTAGCTGAATGTTGTTGAACGAGTTATGTACATCAAAAGA  
TTGAATAAAACAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

**SEQ ID NO:2**  
Amino Acid Sequence  
383 residues  
Mosquito arrestin 1

MVYNFKVFKKCAPNGKVTLYMGKRDFVDHVSVEPIDGIVVLDDEYIRDNRKVF  
GQIVCSFRYGREEDEV MGLNFQKELCLASEQIYPRPEKSDKEQTKLQERLLKKLG  
SNAIPFTFNISP NAPSSVTLQQGEDDNGDPCGVSYVVKIFAGESETDRTHRRSTVT  
LGIRKIQFAPTKQGQQPCTLVRKDFMLSPGELELEVTLDKQLYLHGERIGVNICIR  
NNSNKMVKKIKAMVQQGV DVVLFQNGSYRNTVASLETSEGCPIQPGSSLQKVMY  
LTPLLSSNKQRRGIALDGQIKRQDQCLASTLLAQPDQRDAFGVIISYAVKVKLFL  
GALGGELSAELPFVLMHPKPGTKAKVIHADSQADVETFRQDTIDQQASVDFE

**SEQ ID NO:3**  
cDNA Nucleic Acid Sequence  
1239 nucleotides  
Mosquito odorant receptor 1

ATGAAGCTGAACAACTGAACCCACGGTGGGATGCGTACGATCGACGGGATTCGTTCTGGTTGCAGTTGCTTTGTTT  
GAAATATTTAGGCCTATGGCCACCGGAAGATACGGATCAGGCAACGCGGAACCGGTACATCGCGTACGGTTGGGCTT  
TGCGGATCATGTTTCTACATCTGTACGCTCTAACGCAAGCCCTATACTTCAAGGATGTGAAGGATATTAATGACATC  
GCAAATGCATTGTTTCGTGCTTATGACTCAAGTGACGTTGATCTACAAGCTGGAAAAGTTTAACTACAACATCGCACG  
GATTCAGGCTTGTCTGCGCAAGCTTAACTGCACACTGTATCACCCGAAACAGCGCGAAGAATTCAGCCCCGTTTTAC  
AATCGATGAGTGGAGTGTGTTGGCTGATGATCTTTCTCATGTTTGTGGCTATCTTCACCATCATCATGTGGGTATG  
TCGCCAGCCTTCGACAATGAACGTCGTCTGCCCGTGCCGGCCTGGTTCCCGGTGGACTATCACCATTCTGGACATAGT  
GTACGGTGTACTGTTCTGTATCAAACCATTTGGAATCGTCATGAGCGCAACGTACAACCTCTCGACCGATACCATGT  
TTTCCGGCTTGATGCTACACATAAATGGACAAATTGTGCGGCTTGGTAGTATGGTTAAAAAGCTTGGACATGACGTC  
CCTCCCGAACGCCAATTGGTCGCAACGGATGCGGAATGGAAAGAGATGCGAAAGCGCATCGACCATCACTCCAAAGT  
GTACGGTACGATGTACGCTAAAGTAACGGAGTGTGTGCTGTTTCACAAGGACATCTTAAGGATCTATCTTCGCGCAA  
GTATGCGCGTCTGTAATTATCATTTGTATGACACTGCTGCAACTACCGGGGGCGATGTTACGATGGCCGATCTGCTG  
GGCTGTGGGGTCTATTTGCTAGTAAAGACATCGCAAGTGTTTATTTTCTGTTACGTAGGGAATGAAATCTCCTATAC  
GACGGATAAATTTACAGAGTTTGTGTTGGGTTTTCCAACCTACTTCAAGTTCGATAAGCGTACCAGCCAAGCAATGATAT  
TTTTTCTGCAATGACTCTTAAAGATGTTACATCAAGGTGGGAAGTGTCTTGAAGGTTACGCTAAATCTTCACACA  
TTTTTGCAGATTATGAAGCTATCGTACTCCTATCTGGCCGTACTTCAGAGCATGGAATCAGAGTAATGGTGTTAATA  
TCCTTAA



**SEQ ID NO:4**  
Amino Acid Sequence  
394 residues  
Mosquito odorant receptor 1

5  
MKKDSFFKMLNKHRLCLWPPEDTDQATRNRYLAYGWALRIMFLHLYALTQA  
LYFKDVKDINDIANALFVLMTQVTLIYKLEKFNYNRIARIQACLRKLNCTLYHPKQ  
REEFSPVLQSMGVFWLMIFLMFVAIFTIIMWVMSPAFDNERRLPVPWFVPVDY  
HHSDIVYGVFLYQTIGIVMSATYNFSTD TMFSGLMLHINGQIVRLGSMVKKLG  
10 HDVPPERQLVATDAEWKEMRKRIDHHSKVYGTMYAKVTECVLFHKDILRIYLR  
ASMRVCNYHLYDTAATTGGDVTMADLLGCGVYLLVKTSQVFIFCYVGNEISYTD  
KFTEFVGFSNYFKFDKRTSQAMIFFLQMTLKDVIKVGSVLKVTLNLHTFLQIM  
KLSYSYLA VLQSMES EZ

**SEQ ID NO:5**  
cDNA Nucleic Acid Sequence  
1142 nucleotides  
Mosquito odorant receptor 2

15  
20  
25  
30  
35  
ATGCTGATCGAAGAGTGTCCGATAATTGGTGTCAATGTGCGAGTGTGGCTGTTCTGGTCGTATCTGCGGCGGCCGCG  
GTTGTCCCGCTTTCTGGTCTGGCTGCATCCCGGTGCGCGTGCTGAACGTTTTCCAGTTCCTGAAGCTGTACTCGTCCT  
GGGGCGACATGAGCGAGCTCATCATCAACGATACTTTACCGTGCTGTACTTTAACCTCGTCCTCCGAACCTCCTTT  
CTCGTGATCAATCGACGGAAATTTGAGACATTTTTTGAAGGCGTTGCCGCCGAGTACGCTCTCCTCGAGAAAAATGA  
CGACATCCGACCCGTGCTGGAGCGGTACACACGGCGGGGACGCATGCTATCGATATCGAATCTGTGGCTCGGCGCCT  
TCATTAGTGCTGCTTTGTGACCTATCCTCTGTTTGTGCCCCGGGCGCGGCCTACCGTACGGCGTCACGATACCGGGC  
GTGGACGTGCTGGCCACCCCGACCTACCAGGTCTGTTTGTGCTGCAGGTTTACCTTACCTTCCCCGCCTGCTGCAT  
GTACATCCCGTTTACCAGCTTCTACGCGACCTGCACGCTGTTTGCCTCGTCCAGATAGCGGCCCTAAAGCAACGGC  
30 TCGGACGCTTGGGGCGCCACAGCGGCACGATGGCTTCGACCGGACACAGCGCCGGCACACTGTTCCGCCGAGCTGAAG  
GAGTGTCTAAAGTATCACAAACAAATCATCCAATATGTTTCATGATCTCAACTCACTCGTCACCCATCTGTGTCTGCT  
GGAGTTCCTGTCGTTCCGGATGATGCTGTGCGCACTGCTGTTTCTGCTAAGCATTAGCAATCAGCTGGCACAGATGA  
TAATGATTGGATCGTACATCTTCATGATACTCTCGCAGATGTTTGCCTTCTATTGGCATGCGAACGAGGTACTGGAG  
CAGAGCCTAGGCATTGGCGATGCCATTTACAATGGAGCGTGGCCGGACTTTGAGGAACCGATAAGGAAACGGTTGAT  
35 TCTAATTATTGCACGTGCTCAGCGACCGATGGTGGTAAGATTAAAGTCGGCAACGTGTACCCGATGACGTTGGAAAT  
GTTTCAAAAATTGCTCAACGTGTCCTACTCCTATTTCACTGCTGCGCCGAGTGTACAAC TAA

**SEQ ID NO:6**  
Amino Acid Sequence  
380 residues  
Mosquito odorant receptor 2

40  
MLIEECPIIGVNVRVWLFWSYLRRPRLSRFLVGCIPVAVLNVFQFLKLYSSWGDM  
SELIINGYFTVLYFNLVLRTSFLVINRRKFETFFEGVAAEYALLEKNDDIRPVLER



YTRRGRMLISISNLWLGAFIGACFVITYPLFVPGRGLPYGVITIPGVVDVLATPTYQVV  
FVLQVYLTFPACCMYIPFTSFYATCTLFALVQIAALKQRLGRLGRHSGTMASTGH  
SAGTLFAELKECLKYHKQIIQYVHDLNSLVTHLCLLEFLSFGMMLCALLFLLSIS  
NQLAQMIMIGSYIFMILSQMFIFYWHANEVLEASLGIGDAIYNGAWPDFEPIRK  
5 RLILIIARAQPTDGGKIKVGNVYPMTLEMFQKLLNVSSYSYFTLLRRVYN

**SEQ ID NO:7**  
cDNA Nucleic Acid Sequence  
10 1236 nucleotides  
Mosquito odorant receptor 3

15 ATGCCTTCTGAGCGGCTTCGTCTCATTACTTCCTTCGGAACCTCCTCAAGACAAACGCACGATGGTACTGCCAAAATT  
AAAGGATGAAACAGCAGTGATGCCGTTTCTGCTGCAAATTCAAACCATTGCCGGACTGTGGGGTGACCGTTCCCAGC  
GGTACCGTTTTTATCTCATCTTTTCTACTTCTGCGCGATGGTGGTTCTACCCAAAGTGCTGTTCCGTTATCCAGAT  
CTCGAGGTTGCGGTACGCGGCACGGCCGAGCTGATGTTTGAATCGAACGCATTCTTCGGCATGCTAATGTTTTCTTT  
TCAACGCGACAACCTACGAGCGATTGGTGATCAGCTGCAGGATCTGGCAGCTCTAGTCCTCCAAGACCTACCCACAG  
AGCTGGGAGAGTACCTGATCTCAGTGAACCGACGGGTGATCGGTTCTCCAAAATTTACTGCTGCTGTCACTTTTCC  
ATGGCAACGTTCTTTTGGTTCATGCCCGTCTGGACGACCTATTCGCGCTACTTTGCTGTGCGCAACAGCACGGAACC  
20 GGTCGAGCACGTGTTGCACCTCGAGGAAGAGCTGTACTTCCTGAACATTTCGGAATTCGATGGCGCACTATACGTTTT  
ATGTGGCCATTATGTGGCCACGATCTATACGCTCGGGTTTACCGGTGGCACAAAGCTGCTGACCATTTCAGCAAT  
GTTAAGTACTGTTTCGGCCATGCTGAAGCTCGTTGCACTCCGAATCCACTGTCTAGCGAGAGTAGCGCAAGACCGAGC  
GGAAAAGGAGCTGAACGAGATTATTTCCATGCATCAGCGGGTACTCAACTGCGTGTTCTCTGCTGGAGACGACATTCC  
GCTGGGTATTTTTTCGTGCAGTTCATTGAGTGACAATGATCTGGTGAGTCTCATCTCTACATAGCGGTGACGGGG  
25 TTCAGCTCGACGGTAGCGAATGTATGTGTCCAGATCATTTTGGTGACGGTGGAAACTTACGGCTACGGCTACTTCGG  
AACAGATCTAACCACGGAGGTGCTTTGGAGCTATGGCGTTGCCCTCGCCATTTACGATAGCGAGTGGTACAAGTTTT  
CCATTTTCGATGCGCCGCAAACTTCGACTGCTACTGCAACGATCCCAAAAACCGCTCGGCGTAACGGCGGGAAAGTTT  
CGCTTCGTCAATGTGGCCAGTTTGGCAAGATGCTCAAGATGTCCTATTCATTTTACGTAGTACTGAAGGAGCAGTT  
30 TTAG

**SEQ ID NO:8**  
Amino Acid Sequence  
35 411 residues  
Mosquito odorant receptor 3

MPSERLRLITSFGTPQDKRTMVLPKLKDETAVMPFLLQIQTIAGLWGDRSQRYR  
FYLIFSIFCAMVVLPKVLFGYPDLEVAVRGTAELMFESNAFFGMLMFQFQRDNY  
ERLVHQLQDLAALVLQDLPTLGEYLISVNNRRVDRFSKIYCCCHFSMATFFWFM  
PVWTTYSAYFAVRNSTEPVEHVLHLEEELYFLNIRTSMAHYTFYVAIMWPTIYTL  
40 GFTGGTKLLTIFSNV KYCSAMLKLVALRIHCLARVAQDRAEKELNEIISMHQ RVL  
NCVFLLETFRWVFFVQFIQCTMIWCSLILYIAVTGFSSTVANVCVQIILVTVETY  
GYGYFGTDLTTEVLWSYGVALAIYDSEWYKFSISMRRKLRLLLQRSQKPLGVTA  
GKFRFVNVAQFGKMLKMSYSFYVVLKEQF

**SEQ ID NO:9**  
**Genomic Nucleic Acid Sequence**  
3895 nucleotides  
Mosquito odorant receptor 1

5 AGCTTTGTTTCATTTATGTTGAAATCTAGCCCATTTTGTATAGTGCTGAACGACGAAGAACATACGAAAGTACCTCGT  
CCGAACACTATCAACATTAATTATACCAAGCTAGAAGAAGATATTTATAGTCAAGCCTCAACATCATAGGAACTTT  
AGCAAAACCATTTAATTTACATGATGATAAGTCCCACCTCTTACCCCAGCACAGGTTTGAGAAGGACGAAAGTATCT  
10 TTACGATAATATTACTCTAAGGTAGTTTTTGAATAAAATAAAAATTTACGTGCAAGTGGTGGCATCGGACATCATTC  
GAAAGAATCTACTAAGTCATACACACACCCAAGACGACCGACGTAGTTTCATCTAGAAAAACGGGTCAGCTCCATC  
GAACACGTCAGGACATAACTGCGACATGCGTATGGTCAGTTCCACTAGTGCCAACACTGGTTCAGGGCACTACCTT  
CCGAAGCAGTAGAACCTAATGTATTGGAAATTATTAGGACATACTGCAACATGCATATGGCTAGTTCCGCTGGTACC  
AACGATGGCACCAGGACACTATCTGCGGCCTTGTAATACTACTGTAAATCTATACAAAACGGCTTTACCCATACT  
TTATCACAAAACGGCAGGTGAGGGCTGGATTGCTTCAAAGCATTAGAAATATATAATTTCAAAGTCCATAATCTCC  
15 TTAAAGATAGACAaCAGTAGAGAACACATTTAGTGCTCTTTTCGTTTCGAGTTAGTTGCCTTCTCAAGTAAGCGTTT  
AATGCTCAATTGTTGTAGATTCGTTGGATGACTCTCGCTACGTGCTATAGTGGTCAATACTTCCAATTAGATTTTCAT  
AATTAGTTTCCAATTGTCCACGGAAAACCCaCAAAAGAAAAAAAACCTTGATCTAGGGTGGAATTTTTTCGAGAACA  
ATTGGACACTTCATATGAAAAAGGACAGCTTTTTCAAATGTTAAATAAACACCGTTGGATCCTTTggttgatttca  
attctccaaattctgcagaataattctgcaaattttacaaaactgctcaaccaccaataattccaattaatcatctg  
20 aacattttaaactgataattaagatgagtaattgcttcgctcatcacctaagaaatcgattagtttgataaaaaagaa  
caaattgaaatacaataaagtcacctgaattttattcgaataacggcttgaactcatttatttcaaaaacctttgaga  
aattcctcggtgaaaattggtctcctatagttctgctaacgggcccacttcaaaagcaagaactaacaaaatcataat  
tatgggtgcaagtaactatcagtaggaatcgccattaaaaacttttccctcaatttgcggtcggttaccgggctaaa  
25 tacagagcagagtaacgggaagtgatcaacgctcgctattagtagtaacgaggaacgcccctccgaagggtgtgttgagg  
accttttcaaattgaaaccaagtactgtttccagtttttaattggatagttataaaatgagccggtcaacgatcggg  
catcatttgagtttcatcttcgaggagaaatagatcagtgccactgtttaaccgaaagtaataagctgaacaaact  
gaaccacgggtgggatgctgacgatcgacgggattcggtctggttgagttgctttggttgaaatatttagGCCTAT  
GGCCACCGGAAGATACGGATCAGGCAACGCGGAACCGGTACATCGCGTACGGTTGGGCTTTGCGGATCATGTTTCTA  
CATCTGTACGCTCTAACGCAAGCCCTATACTTCAAGgATGTGAAGGATATTAATgtgagttcttagtttagctattag  
30 tgttccacctgtccataatctgtcttttattgggtagGACATCGCAAATGCATTGTTTCGTGCTTATGACTCAAGTGA  
CGTTGATCTACAAGCTGGAAAAGTTTAACTACAACATCGCACGGATTACAGGCTTGTCTGCGCAAGCTTAACTGCACA  
CTGTATCACCCGAAACAGCGCGAAGAATTCAGgtaagcctgctgggaaatatgactaaaaagagtgttaacaaacga  
ctctcctccaaatgtagCCCCGTTTTACAATCGATGAGTGGAGTGTGTTGGCTGATGATCTTTCTCATGTTTGTGGC  
TATCTTCACCATCATCATGTGGGTTATGTGCGCCAGCCTTCGACAATGAACGTCGTCTGCCcGTGCCGGCCTGGTTCC  
35 CGGTGGACTATCACCATTCCGACATAGTGTACGGTGTACTGTTCTGTATCAAACCATTGGAATCGTCATGAGCGCA  
ACGTACAACCTTCTCGACCGATAACCATGTTTTCCGGCTTGATGCTACACATAAATGGACAAATTGTGCGGCTTGGTAG  
TATGGTTAAAAAGgtgagttacggcgactacttgctccagtaaggacagggagtttggttccggttatgatatcatt  
ttatcagCTTGGACATGACGTCCCTCCCGAACGCCAATTGGTTCGCAACGGATGCGGAATGGAAAGAGATGCGAAAGC  
GCATCGACCATCACTCCAAAGTGACGGTACGATGTACGCTAAAGTAACGGAGTGTGTGCTGTTTCACAAGGACATC  
40 TTAAGgtacgaattgggccaattaattgtgtcattttaaagcttgacccaacttttcacagcttcggcgatgaagt  
gcaggacattttccaagGATCTATCTTCGCGCAAGTATGCGCGTCTGTAATTATCATTTGTATGACACTGCTGCAAc  
TACCGGGGgCGATGTTACGATgCCCGATCTGCTGGGCTGTGGGGTCTATTTGCTAGTAAaGACATCGCAAGTGTTTA  
TTTTCTGTTACGTAGGGAATGAAATCTCCTATACGgtaggttgagacacgtagaggaattaaatgtttggaagaata  
tcaataccaaatagtatgatgtttcggttacagACGGATAAATTTACAGAGTTTGTGTTGGGTTTTCCAACACTACTTCAAG  
45 TTCGATAAGCGTACCAGCCAAGCAATGATATTTTTTCTGCAAATgtgagatagcgggtgtatttgtgcagtcagtaca  
ttaaatacgttctctattttcagGACTCTTAAAGATGTTTACATCAAGGTGGGAAGTGTCTTGAAGGTTACGCTAAAT  
CTTCACACATTTTTGTCAGgtatgtaattatgctgtggtatttagcttgaaataagctacaaactttgaaagtaattt  
caatctgtttttagATTATGAAGCTATCGTACTCCTATCTGGCCGTACTTCAGAGCATGGAATCAGAGTAATGGtG  
tTAATATCCtTAATGTTGAAATTATATTTTGTAGATTTATTGCATAAAGTAaTaTTTAATTTTATACATCAAACGT  
50 AAGCCCGctaGTTTTCAATTAGCCTTTTTCCAAAATTTATCAAATTGATTTTGAATTGATTGCAGAGTTTCAGGAATT  
TAATCTGATAGGATATCTTGTATTATCCAATAGAGGTGTGGAAGCGTTCCCAAGCCATTCGTTTGATAGTTTATAGCA

CCGTCGAGCAGTTGATCGCTGTGATCGCTAGGCGCACCTGATTTTATCTTTATCTCGCACCTGTTATGGCAAGGGCG  
CTTTTCACACGTTTTCACACAATATAATGCACATGTATAATGCATTCTTACTTTAGCATTTTTGTTACATATAATACC  
AAAATTATGCATTTTTTATTCTCACGCAACGATTAGAGGATGACTTcACAAAGGTCCATCTAGTGGTAGGAGGTATAC  
AATTATACCTCTCAAATCTCACAGCAtAATGAGAAACAAAAGGATACCAAGCATAACCCTTTTTTTTACTTGACAATT  
5 TCATTTGATTTATGTAATAAAGCACTGCaCGTCGACTTCCTAAAA

**SEQ ID NO:10**

10 Genomic Nucleic Acid Sequence  
4985 nucleotides  
Mosquito odorant receptor 2

15 GGGATCCTCTAGAGTCGACCTGCAGGCATGCAAGCTTCCCTCACCGTGACGTGCTAGAAATGGTTCAACATACTCGT  
CCGGCAGAGCGAAGACGACGAACAGCGGAATGTCCCAGGAAATGTAATGAGATATCACAGCAAGTGAACCCAAACCG  
AGCTGTGCGCTTTGTGTTGCGCTTTAAAAATGGCCCTTCCTTCGCCGCATCTGCTTGGTTTCACACGCTTTCCAGG  
AAATCCACTGACCACTGGCCACACATCAACCACCGGAGCGGGAGCCTCAGTGCCAGCGAAGCATATAATTTGCTCA  
AAAAGTCACGGTACTCAATTAATTTGATTATAATCAATTTTCGTGGCTTCCAACACACCCTTCTTCCACAATCCATCG  
20 CCGAGTGAGCGAGTATAAAGGTGAAGAAACGTACCTTGCGCTTGCTCACTAACTGAACCGGATTTCAAAAAGGAACA  
TAAACCGCAACCCACAGCCGAAAATGCTGATCGAAGAGTGTCCGATAATTGGTGTCAATGTGCGAGTGTGGCTGTTC  
TGGTCGTATCTGCGGCGGCCGCGGTTGTCCCGCTTTCTGGTCGGCTGCATCCCGGTGCGCGTGCTGAACGTTTTCCA  
GTTCTGAAGCTGTACTCGTCCTGGGGCGACATGAGCGAGCTCATCATCAACGGATACTTTACCGTGCTGTACTTTA  
ACCTCGTCGTacgtgggaggggaggggcaataaccttcccacttggtggatattttcataccttttccatgtggt  
35 tttttatttctctgtttgttgccatccagCTCCGAACCTCCTTTCTCGTGATCAATCGACGGAAATTTGAGACATTTT  
TTGAAGGCGTTGCCGCCGAGTACGCTCTCCTCGAGgtaagtcattgggtttttctagtttttgggggagttggtttaca  
ccataaccacccccgacggtaacatttgatcgtcccgcgaaaatgtttgtagagAAAAATGACGACATCCGACCCGT  
GCTGGAGCGGTACACACGGCGGGGACGCATGCTATCGATATCGAATCTGTGGCTCGGCGCCTTCATTAGTGCCTGCT  
TTGTGACCTATCCTCTGTTTGTGCCCCGGGCGCGGCCTACCGTACGGCGTCACGATACCGGGCGTGACGTGCTGGCC  
ACCCCGACCTACCAGGTCGTGTTTGTGCTGCAGGTTTACCTTACCTTCCCCGCCTGCTGCATGTACATCCCGTTCAC  
40 CAGCTTCTACGCGACCTGCACGCTGTTTGTGCTCGTCCAGATAGCGGCCCTAAAGCAACGGCTCGGACGCTTGGGGC  
GCCACAGCGGCACGATGGCTTCGACCGGACACAGCGCCGGCACACTGTTTCGCCGAGCTGAAGGAGTGTCTAAAGTAT  
CACAAACAAATCATCCAGtaagtagacgctagtagactcgaccggattgcccttccttcggggaggggaggtttgct  
atttcgggatgcggcagcacgcatacacacaaaccggaagccattaattctcccgttttcatgcccgcacgggcact  
gggtcatgtttcacatccttccttcctttccaaacacacacacgcgcgcgtgcacgtacagATATGTTTCATGATCTC  
50 AACTCACTCGTCACCCATCTGTGTCTGCTGGAGTTCCTGTCTGTTTCGGGATGATGCTGTGCGCACTGCTGTTTCTGCT  
AAGCATTGtaagtaaaatcgaccgacgtgcggtcgctagtcctcctcggaactctcatttcgggactcaatcgttcc  
atctctcaatagAGCAATCAGCTGGCACAGATGATAATGATTGGATCGTACATCTTCATGATACTCTCGCAGATGTT  
TGCTTCTATTGGCATGCGAACGAGGTACTGGAGCAGGtaatggcgctgaagctgagtttggttgagcgggttcgcta  
tagatcggctgtcttacattggttggtttctgcacgtggggatcggttttggttttctctccatttcagAGCCTAGGC  
ATTGGCGATGCCATTTACAATGGAGCGTGGCCGGACTTTGAGGAACCGATAAGGAAACGGTTGATTCTAATTATTGC  
ACGTGCTCAGCGACCGATGGTGGTAAGtttggtgatcgatgctctgttcaatgaacatggcacagaaggctgtgta  
aatagctgttcattaataagttttttcagaatgtatcggttttagttgatttaaacgcattgttctatgcaatggta  
gcaacaatagaccgcctttattaatccaagcttcctttaggattgattttttatttttaagagaaagataaaccatttt  
tagtaaccaatttagttacaggaaccaaatacagaatttattattattattattattattattattattattattatt  
45 ttatt  
att  
tatt  
ttattattattattattattattataacaataataattattattattattattattattattattattattattatt  
att  
50 ttatt

[illegible]

10

15

20

25

520

**SEQ ID NO:11**

Genomic Nucleic Acid Sequence  
2083 nucleotides

### Mosquito odorant receptor 3

AAGCAGAACACATCAAGAAGCAATTAGGTGTGTCGTACGTTAGCAAGTAGTTCGCGAGGAGGAATAAAATAGATGCC  
TTCTGAGCGGCTTCGTCTCATTACTTCCTTCGGAACCTCCTCAAGACAAACGCACGATGGTACTGCCAAAATTAAAGG  
ATGAAACAGCAGTGATGCCGTTTCTGCTGCAAATTCAAACCATTGCCGGACTGTGGGGTGACCGTTCCAGCGGTAC  
CGTTTTTATCTCATCTTTTCCTACTTCTGCGCGATGGTGGTTCTACCCAAAGTGCTGTTTCGGTTATCCAGATCTCGA  
GGTTGCGGTACGCGGCACGGCCGAGCTGATGTTCGAATCGAACGCATTCTTCGGCATGCTAATGTTTTCTTTCAAC  
GCGACAACTACGAGCGATTGGTGCATCAGCTGCAGGATCTGGCAGCTCTAGgtgagtatgagccaatcgattgttc  
caaaccttcgcaacatccttcgtaacactgctacactttcagTCCTCCAAGACCTACCCACAGAGCTGGGAGAGTAC  
CTGATCTCAGTGAACCGACGGGTCGATCGGTTCTCCAAAATTTACTGCTGCTGTCACTTTTCCATGGCAACGTTCTT  
TTGGTTCATGCCCCGTCTGGACGACCTATTCCGCCTACTTTGCTGTGCGCAACAGCACGGAACCGGTCGAGCACGTGT  
TGCACCTCGAGGAAGAGCTGTACTTCCTGAACATTCCGACTTCGATGGCGCACTATACGTTTTATGTGGCCATTATG  
TGGCCACGATCTATACGCTCGGGTTTACCGGTGGCACAAAGCTGCTGACCATTTTCAGCAATGTTAAGTACTGTTC  
GGCCATGCTGAAGCTCGTTGCACTCCGAATCCACTGTCTAGCGAGAGTAGCGCAAGACCGAGCGGAAAAGGAGCTGA  
ACGAGATTATTTCCATGCATCAGCGGGTACTCAAgtaaagtaaattcaaattgaaagttttgcaggggaataaacttgag  
tgtgtctgacccgtgcacatcctagCTGCGTGTTCTTGCTGGAGACGACATTCCGCTGGGTATTTTTTCGTGCAGTTC

ATTCAGTGTACAATGATCTGGTGCAGTCTCATCCTCTACATAGCGGTGACGgtaatagcatttttcgtcattttcgтта  
gccttattcaatccattttttgtgaacgtgaatttccccccagGGTTCAGCTCGACGGTAGCGAATGTATGTGTCCAG  
ATCATTTTGGTGACGGTGGAACTTACGGCTACGGCTACTTCGGAACAGATCTAACCACGGAGGTGCTTTGGgtacc  
ctttggatgaagcttcaaaaagtaattccaaattctgttttcgattttttcccccttttccactagAGCTATGGCGTTG  
5 CCCTCGCCATTTACGATAGCGAGTGGTACAAGTTTTCATTTTCGATGCGCCGCAAACCTTCGACTGCTACTGCAACGA  
TCCCCAAAACCGCTCGGCGTAACGGCGGGAAAGTTTCGCTTCGTCAATGTGGCCCAGTTTGGCAAGgtaacattaat  
tacagtttgaaaattctgaagaatgcatcttacttgcttacttggtgttccagATGCTCAAGATGTCCTATTTCATT  
TTACGTAGTACTGAAGGAGCAGTTTTAGGAGCTGCTGTTTCCACCCCTGGAAATGGCCTTTTCGCACTGTCTTCTGT  
TTGTTGGACGCACGCAGCACCAGAGCGCCCTGCACGCACTGACGTATTTTGGCTACTTTGACGTTTGCACCTTTG  
10 ACAGCTGAAGGACAGGGTACAATTTTTGCTGCTGTTATTACGCGCAGCGCATTGGATACGAAAACATTGGCCACAAG  
TTCTACGATTTTACGCTTTTATTACTGTTTCGTAGCAGCTTTTTTCCaCAATAAACACACACAATAACGTACCGACAG  
TATTCTTTTCATTGTAGGATAGAGAAGCCGCCGCCAGCAGCCAAAACGCGCCGCAAAACGAAAGGCGGCACCACCG  
GGGGAAAACACGGGAGCAAAACGAGAACAGAACGCAGTAAACAACAAAACCGGCCGGAACAACAACGGTGCCGGAA  
ACGA

15

10056405  
2024-07-09 10:56:40

**SEQ ID NO:12**  
Genomic Nucleic Acid Sequence  
2374 nucleotides  
Mosquito odorant receptor 4

GGGGAACTCCCCCACCAGACGACGGAAAGCTAACGATGTGCAATTGAA  
TAGTCATTAGTAGCGTTTTTGTCTCGCAAACGAACTAACCTTTGACTTTTTTAAG  
25 TTCACTACGGTGAGGACAAAAATCAATAAATTAAATCGAGACCGTTGATGAGCA  
AAAGAAAAAAAAAATATTTTACTGATTTTTCATTTTCGTTCCATCGACTACATAATCA  
TAATTATATGCCACATTTTATTATAAGTTTTTGTATCATTTTTTAAACAACACAAA  
AATGCATCCTTTTCGAATATTAGTCAGGTTGTATCAACAATGAAGTTTGAAGTGT  
TTCAAAAATATTCCTCCCCGGACACGGTCTTATCCTTCGTGCTAAGGCTTTTGC  
30 ATATCGTGGGCATGAATGGGGCAGGATTTTCGGTCGCGAATTCGAGTTGGTGGC  
ATTTTTCTGTTCTATTTAATCTTTCTTGTAATACCGCCACTAACGGGCGGGTAC  
ACCGATGGTCACCAGCGTGTACGCACCAGTGTGGAATTCCTGTTTAATTGCAAT  
ATTTACGGCGGCAGTATGTTCTTTGCCTACGATGTGGCCACTTTCCAAGCGTTC  
ATCCAGGAACCTGAAGAGCCTTTTCGGTTTTTGGgtaatatttaattaataaaattgcgtttattgca  
35 tcatcatttgtttctctttgcagTATGCTCACATTCGTACAGACTAAAGTATAAGCTGACCCG  
GTTCAACCGTCGAGCGGATATTATCGCCAAAGTGCAAACGACCTGCATGGGTG  
CTGTAACGCTTTTCTACTGGATTGCACCGATACCTTCCATCTGTGCGCACTACT  
ACAGGTGACCAATTCCACCGAACCCGTGCGGTTTGTGCAACATTTAGAGGTG  
AAGTTCTATTGGCTCGAGAATCGCACCTCAGTCGAGGACTACATAACCTTCGTG  
40 CTGATCATGCTACCCGTCGTGGTTATGTGTGGTTACGTATGCAATTTGAAGGTG  
ATGACCATCTGCTGCAGCATTGGACACTGTACACTGTACACCAGGATGACTATA  
GAGATGGTAGAGCAGTTGGAAAGCATGGCATCAGCGGAACGAACTGCCAGCGC  
CATACGCAACGTGGGGCAGATGCACAGTGGTTTACTGAAATGCATTAGGCTTT  
TGAACACGTCAATCCGATCGATGCTGATGCTGCAGTGGTTGACCTGCGTGTTA  
45 AACTGGAGCATTTCTCTCATCTATCTAACGAACGTGgtagttttgtcttggttgaaatccaa



aaacaaaaagatggctataattgaactttctattacagGGCATCTCGCTACAATCGGTTACCGTGG  
TGGTAATGTTTTTTCTTGCCACTGCGGAACTTTCCTGTATTGTTTACTTGGGA  
CGCGGCTTGCGACACAACAGCAGCTGCTGGAGCACGCACTCTATGCTACACGG  
TGGTACAAC TACCCAATAGCCTTTCGCAGCAGCATTAGGATGATGTTGAGACA  
5 GTCGCAAAGGCATGCACACATAACGGTGGGGGAAGTTTTTTTCGCGTTAATTTGG  
AAGAATTTAGCAGGATTGTCAACTTATCCTACTCTGCTTACGTCGTACTTAAGG  
ATGTAATAAAGATGGATGTACAGTGAATGTTTTTTTTTTTTGGCTTGGCAACGAA  
TGAAGTTTTCCGAATCTATATTAGATCTAGAATTTAATCTAGATGTCATAATATG  
ATCTTGGCCATGACCGGTTCTTGGTTTTGGAACCAATTCTCAAAAACAATTTTGA  
10 ACTTAGGGGCGAGGCATGAAATGTCCCAAGAACCTATCCAAGTTCTGGAAC TAC  
ATATTACCGAATCTATCCCATTATTGCCTCGGAACTGGTTTGGTGCTAAATATT  
TGTCCAAATGTTGGTCCTGGACCTATCCAGACAAAGATCTTCAATTATTCCTAC  
CACTGGAAC TGAATTAATTGATGTAGGAAGTCATGGAGGTGTTTCAGGGAGAATT  
TAAACACTAATGTTCCAAC TCAATTATTTCAAGGGCAATTCTATTTTTTTATATGCC  
15 CCTACGGATTGATACGTATGTATTACTCCATTTCTGACTTTGTCTTATTCTTG  
CTGCTGATTGGACGTGAAATGTTGAGAAAAAGATTCTTATTTATGAGTGATACA  
GAGCCTTTAAATACTCCTACGTTGTTTGCTATTTAAGTATGGCCAGGCTAATCA  
CAATCGCTACTAATGAACAGAATCTCTTCTAATTAAACCCTTTTCGATTGATAGT  
GTCAATGTCAATGTTCGAGATAATTGAACTGCAAACgATACCTACCTTAAACGGA  
20 GCAGAACACATCAAGAAGCAATTAGGTGTGTCGTACGTTAGCAAGTAGTTTCGC  
GAGGAGGAATAAAATAG

**SEQ ID NO:13**

cDNA Nucleic Acid Sequence

1194 nucleotides

Mosquito odorant receptor 4

30 ATGAAGTTTGAAC TGTTCAAAAATATTCCTCCCCGGACACGGTCTTATCCTTCGTGCTAAGGCTTTTGCATATCGT  
GGGCATGAATGGGGCAGGATTTTCGGTCGCGAATTCGAGTTGGTGGCATTTTTCTGTTCTATTTAATCTTTCTTGTA  
TACCGCCACTAACGGGCGGGTACACCGATGGTCACCAGCGTGACGCACCAGTGTTGAATTCCTGTTTAATTGCAAT  
35 ATTTACGGCGGCAGTATGTTCTTTGCCTACGATGTGGCCACTTTCCAAGCGTTCATCCAGGAACTGAAGAGCCTTTC  
GGTTTTGGTATGCTCACATTCGTACAGACTAAAGTATAAGCTGACCCGGTTCAACCGTCGAGCGGATATTATCGCCA  
AAGTGCAAACGACCTGCATGGGTGCTGTAACGCTTTTCTACTGGATTGCACCGATACCTTCCATCTGTGCGCACTAC  
TACAGGTCGACCAATTCACCCGAACCCGTGCGGTTTGTGCAACATTTAGAGGTGAAGTTCTATTGGCTCGAGAATCG  
CACCTCAGTCGAGGACTACATAACCTTCGTGCTGATCATGCTACCCGTCGTGGTTATGTGTGGTTACGTATGCAATT  
TGAAGGTGATGACCATCTGCTGCAGCATTGGGACACTGTACACTGTACACCAGGATGACTATAGAGATGGTAGAGCAG  
40 TTGGAAAGCATGGCATCAGCGGAACGAAC TGCCAGCGCCATACGCAACGTGGGGCAGATGCACAGTGGTTTACTGAA  
ATGCATTAGGCTTTTGAACACGTCAATCCGATCGATGCTGATGCTGCAGTGGTTGACCTGCGTGTTAAACTGGAGCA  
TTTCTCTCATCTATCTAACGAACGTGGGCATCTCGCTACAATCGGTTACCGTGGTGGTAATGTTTTTTCTTGCCACT  
GCGGAACTTTCTGTATTGTTTACTTGGGACGCGGCTTGCGACACAACAGCAGCTGCTGGAGCACGCACTCTATGC  
TACACGGTGGTACAAC TACCCAATAGCCTTTCGCAGCAGCATTAGGATGATGTTGAGACAGTCGCAAAGGCATGCAC

ACATAACGGTGGGGAAGTTTTTTCGCGTTAATTTGGAAGAATTTAGCAGGATTGTCAACTTATCCTACTCTGCTTAC  
GTCGTACTTAAGGATGTAATAAAGATGGATGTACAGTGA

5

**SEQ ID NO:14**

Amino Acid Sequence

412 residues

Mosquito odorant receptor 4

10

MKFELFQKYSSPDTVLSFVLRLLHIVGMNGAGFRSRIRVGGIFLFYLIFLVIPPLTGGYTDGHQRVRTSVEFL  
FNCNIYGGSMFFAYDVATFQAFIQELKSLSVLVCSHSYRLKYKLTRFNRRADIIAKVQTTCMGAVTLFYWI  
APIPSICAHYYRSTNSTEPVRFVQHLEVKFYWLENRTSVEDYITFVLIMLPVVVMCGYVCNLKVMTICCSIG  
HCTLYTRMTIEMVEQLESMAAERTASAIRNVGQMHSGLLKCI RLLNTSIRSMLMLQWLTCVLNWSISLIY  
15 LTNVGISLQSVTVVVMFFLATAETFLYCLLGTRLATQQQLLEHALYATRWNYPPIAFRSSIRMMLRQSQRH  
AHITVGKFFRVNLEEF SRIVNLSYSAYVVLKDVIKMDVQNVSYSYFTLLRRVYN

15

20

**SEQ ID NO:15**

cDNA Nucleic Acid Sequence

1176 nucleotides

Mosquito odorant receptor 5

25

ATGGTGCTACCGAAGCTGTCCGAACCGTACGCCGTGATGCCGCTTCTACTACGCCTGCAGCG  
TTTCGTTGGGCTGTGGGGTGAACGACGCTATCGCTACAAGTTCGGGTTGGCATTTTTAAGCTT  
CTGTCTGCTAGTAGTTATTCCGAAGGTTGCCTTCGGCTATCCAGATTTAGAGACAATGGTTCG  
CGGAACAGCTGAGCTGATTTTCGAATGGAACGTACTGTTTGGGATGTTGCTGTTTTCTCTCAA  
GCTAGACGACTATGATGATCTGGTGTACCGGTACAAGGACATATCAAAGATTGCTTTCCGTA  
30 AGGACGTTCCCTCGCAGATGGGCGACTATCTGGTACGCATCAATCATCGTATCGATCGGTTT  
TCCAAGATCTACTGCTGCAGCCATCTGTGTTTGGCCATCTTCTACTGGGTGGCTCCTTCGTCC  
AGCACCTACCTAGCGTACCTGGGGGCACGAAACAGATCCGTCGCCGGTCGAACATGTGCTAC  
ACCTGGAGGAGGAGCTGTACTGGTTTCACACCCGCGTCTCGCTGGTAGATTACTCCATATTC  
ACCGCCATCATGCTGCCTACAATCTTTATGCTAGCGTACTTCGGTGGACTAAAGCTGCTAAC  
35 CATCTTCAGCAACGTGAAGTACTGTTTCGGCAATGCTCAGGCTTGTGGCGATGAGAATCCAGT  
TCATGGACCGGCTGGACGAGCGCGAAGCGGAAAAGGAACTGATCGAAATCATCGTCATGCA  
TCAGAAGGCGCTAAAATGTGTGGAGCTGTTGGAAATCATCTTTCGGTGGGTTTTTCTGGGAC  
AGTTCATACAGTGCCTAATGATCTGGTGCAGCTTGGTTCTGTACGTCGCCGTTACGGGTCTCA  
GCACAAAAGCGGCAAACGTGGGTGTACTGTTTATACTGCTAACAGTGGAACCTACGGATTTC  
40 TGCTACTTTGGCAGTGATCTTACCTCGGAGGCAAGTTGTTATTCGCTGACACGTGCTGCGTAC  
GGTAGCCTCTGGTATCGCCGTTTCGGTTTCGATTCAACGGAAGCTTCGAATGGTACTGCAGCG  
TGCCCAGAAACCGGTCGGCATCTCGGCTGGGAAGTTTTGCTTCGTCGACATTGAGCAGTTTG  
GCAATATGGCAAAAACATCATACTCGTTCTACATCGTTCTGAAGGATCAATTTTAA

45

**SEQ ID NO:16**

Amino Acid Sequence

391 residues

5 Mosquito odorant receptor 5

MVLPKLSEPYAVMPLLLRLQRFVGLWGERRYRYKFRLAFLSFCLLVVIPKVAFGYPDLETMVRGTAE LIFE  
WNVLFGLMLLSLKLDDYDDL VYRYKDISKIAFRKDVPSQMGDYLVRINHRIDRFSKIYCCSHLCLAIFYWV  
APSSSTYLAYLGARNRSVPVEHVLHLEEELYWFHTRVSLVDYSIFTAIMLPTIFMLAYFGGLKLLTIFS NVK  
10 YCSAMLRLVAMRIQFMDRLDEREAEKELIEIIVMHQKALKCVELLEIIFRWVFLGQFIQCVMIWCSLVLYVA  
VTGLSTKAANVGVLFI LLTVETYGFCYFGSDLTSEASCYSLTRAAYGSLWYRRSVSIQRKLRMVLQRAQKP  
VGISAGKFCFVDIEQFGNMAKTSYSFYIVLKDQF

**SEQ ID NO:17**

Partial cDNA Nucleic Acid Sequence

474 nucleotides

Mosquito odorant receptor 6

TTATGCTTACCGGATGTTGCGATCGCGCACGTGCTTTTCCGCATACGCCAGTGCACACTTGAT  
GGCGGTGGTGATGACGTCTGCTGCGCACCGTTTTCTGCTCGTGAGTCAGACCTTTTCATTTC  
TGCAATATCCTGTTTCTTTCCCGACCCACAGACGGTTAGACGGATATATGCTGGTAAAGTTT  
GTCCTCTTCATGCTGTGCTTTCTGATCGAGCTGCTGATGCTGTGTGCGTACGGTGAGGATATT  
20 GTGGAATCGCCTTGGGGTGATTGATGCCGCTTACGGTTGCGAATGGTACCGGGAAGGGTCGG  
TGGCGTTCCATCGATCCGTGCTGCAAATTATACACCGCAGCCAGCAGTCCGTCATACTGACC  
GCATGGAAAATTTGGCCCATCCAAATGAGTACTTTCAGTCAGATCCTGCAAGCTTCCTGGTC  
CTACTTTACCCTCCTGAAGACCGTCTACGGGAATAA

**SEQ ID NO:18**

Partial Amino Acid Sequence

157 residues

35 Mosquito odorant receptor 6

LCLPDVAIAHVLFRIRQCTLDGGGDDVCCAPFSARESDFISCNILFLSRPHRRLDGYMLVKFVLFMLCFLIE  
LLMLCAYGEDIVESPWGDZCRLRLRMVPGRVGGVPSIRAANYTPQPAVRHTDRMENLAHPNEYFQSDPAS  
FLVLLYPPEDRLRE

**SEQ ID NO:19**

cDNA Nucleic Acid Sequence



1206 nucleotides

Mosquito odorant receptor 7

5 ATGGTGCTGATCCAGTTCTTCGCCATCCTCGGCAACCTGGCGACGAACGCGGACGACGTGAA  
CGAGCTGACCGCCAACACGATCACGACCCTGTTCTTCACGCACTCGGTACCAAGTTCATCT  
ACTTTGCGGTCAACTCGGAGAACTTCTACCGGACGCTCGCCATCTGGAACCAAGACCAACACG  
CACCCGCTGTTTGCCGAATCGGACGCCCAGTACCATTCGATTGCGCTCGCCAAGATGCGGAA  
GCTGCTGGTGCTGGTGATGGCCACCACCGTCCTGTCGGTTGTCGCCTGGGTACGATAACAT  
TTTTCGGCGAGAGCGTCAAGACTGTGCTCGATAAGGCAACCAACGAGACGTACACGGTGGA  
10 TATACCCCGGCTGCCCATCAAGTCCTGGTATCCGTGGAATGCAATGAGCGGACCGGCGGTACA  
TTTTCTCTTTCATCTACCAGGTACGTTGGCGGAATGGTATTATGCGATCGTTGATGGAGCTTT  
CGGCCTCGCTGGACACCTACCGGCCCAACTCTTCGCAACTGTTCCGAGCAATTTAGCCGGT  
TCCAAATCGGAGCTGATCATCAACGAAGAAAAGGATCCGGACGTTAAGGACTTTGATCTGA  
GCGGCATCTACAGCTCGAAGGCGGACTGGGGCGCCCAGTTCCGTGCGCCGTCGACGCTGCA  
15 AACGTTTCGACGAGAATGGCAGGAACGGAAATCCGAACGGGGCTTACCCGGAAGCAGGAAAT  
GATGGTGCGCAGCGCCATCAAGTACTGGGTCGAGCGGCACAAGCACGTTGTACGTCTCGTTT  
CAGCAATCGGAGATACGTACGGTCCTGCCCTGCTGCTACACATGCTGACCTCCACCATCAAG  
CTGACGCTGCTCGCCTACCAGGCAACGAAAATCGACGGTGTCAACGTGTACGGATTGACCGT  
AATCGGATATTTGTGCTACGCGTTGGCTCAGGTTTTCTGTTTTGCATCTTTGGCAATCGGCT  
20 CATCGAGGAGAGCTCATCCGTGATGAAGGCGGCCTATTCCTGCCACTGGTACGACGGGTCCG  
AGGAGGCAAAAACCTTCGTCCAGATCGTTTGTGAGCAGTGCCAGAAGGCGATGACTATTTCC  
GGAGCCAAGTTTTTCACCGTTTCGCTCGATCTGTTTGCTTCGGTTCCTGGAGCCGTTGTCACC  
TACTTCATGGTGCTGGTGCAGCTGAAGTAA

**SEQ ID NO:20**

Amino Acid Sequence

401 residues

30 Mosquito odorant receptor 7

MVLIQFFAILGNLATNADDVNELTANTITTLFFTHSVTKFIYFAVNSENFYRTLAIWNQTNTHPLFAESDAR  
YHSIALAKMRKLLVLVMATTVLSVVAWVTITFFGESVKTVLDKATNETYTVDIPRLPIKSWYPWNAMSGP  
AYIFSFIYQVRWRNGIMRSLMELSASLDTYRPNSSQLFRAISAGSKSELINEEKDPDVKDFDLSGIYSSKAD  
35 WGAQFRAPSTLQTFDENGRNGNPNGLTRKQEMMVRSIAIKYWVERHKKHVRLVSAIGDTYGPALLLHMLT  
STIKLTLLAYQATKIDGVNVYGLTVIGYLCYALAQVFLFCIFGNRLIESSSVMKAAYSCHWYDGSEEAKTF  
VQIVCQQCQKAMTISGAKFFTSLDLFASVLGAVVTYFMVLVQLK

**SEQ ID NO:21**

Genomic Nucleic Acid Sequence

2272 nucleotides

Mosquito odorant receptor 5

5 tctagactgaacccatgacgggcattttattgagtcgttcgagttgacgactgtaccacgggaccacccgtttatcactatcactattaattaattataatgc  
ttttagcgcgatcagcctaccgggtttgtttctctggatatcttaagttccatttgattatcaagatagaacaacaactgtaccttaataatcattacgtacc  
ttaatcaacctgtgcatcaaggagtttgcgaaagcaaaaatccgattgtctgatgttcttgattccatccgattcgttactggttctgcaaaatcgtccat  
aatacggcaatgtccttatcgcgtgctgaatcaacatcacattgtttgcatttcgtttttgcgtgcaaatatgttattgcaaagaaggcaaggtaatgtgetta  
10 gagtaaatacaattcgcgtgctccattttgtccaccagtgtgccagaacccgtgccttttagtccttcgaatacatccgaccagtgcgaagcaagtgcacA  
TGGTGCTACCGAAGCTGTCCGAACCGTACGCCGTGATGCCGCTTCTACTACGCCTGCAGCGT  
TTCGTTGGGCTGTGGGGTGAACGACGCTATCGCTACAAGTTCCGGTTGGCATTTTTAAGCTTC  
TGTCTGCTAGTAGTTATTCCGAAGGTTGCCTTCGGCTATCCAGATTTAGAGACAATGGTTTCG  
GGAACAGCTGAGCTGATTTTCGAATGGAACGTACTGTTTGGGATGTTGCTGTTTTCTCTCAAG  
15 CTAGACGACTATGATGATCTGGTGTACCGGTACAAGGACATATCAAAGATTGgtgcgtgataatgattg  
ataaaaggaacctttgagcaactcctatcccttcaagCTTTCCGTAAGGACGTTCCCTCGCAGATGGGCGACTATCT  
GGTACGCATCAATCATCGTATCGATCGGTTTTCCAAGATCTACTGCTGCAGCCATCTGTGTTT  
GGCCATCTTCTACTGGGTGGCTCCTTCGTCCAGCACCTACCTAGCGTACCTGGGGGACGAA  
ACAGATCCGTCCTCGGTGGAACATGTGCTACACCTGGAGGAGGAGCTGTACTGGTTTCACACC  
20 CGCGTCTCGCTGGTAGATTACTCCATATTCACCGCCATCATGCTGCCTACAATCTTTATGCTA  
GCGTACTTCGGTGGACTAAAGCTGCTAACCATCTTCAGCAACGTGAAGTACTGTTTCGGCAAT  
GCTCAGGCTTGTGGCGATGAGAATCCAGTTCATGGACCGGCTGGACGAGCGCGAAGCGGAA  
AAGGAACTGATCGAAATCATCGTCATGCATCAGAAGGCGCTAAAgtaaggctgcccgtatgtgtggataga  
atacatttctagctgctttcagATGTGTGGAGCTGTTGGAAATCATCTTTCGGTGGGTTTTCTGGGACAGT  
25 TCATACAGTGCGTAATGATCTGGTGCAGCTTGGTTCTGTACGTCGCCGTTACGgtaactaaaagcactg  
tagtgatctgtctgccacaccattcactgctgtgtcttgtttgtcactctccagGGTCTCAGCACAAAAGCGGCAAACGTGGG  
TGTACTGTTTATACTGCTAACAGTGGAACCTACGGATTCTGCTACTTTGGCAGTGATCTTAC  
CTCGGAGGCAAGTTGTTATTCGCTGAgtttcagttactttccgttccctctaaccgtaccattgtaccattgttgagacagagc  
ttgagcgtagCACGTGCTGCGTACGGTAGCCTCTGGTATCGCCGTTTCGGTTTCGATTCAACGGAAG  
30 CTTCGAATGGTACTGCAGCGTGCCAGAAACCGGTCGGCATCTCGGCTGGGAAGTTTTGCTT  
CGTCGACATTGAGCAGTTTGGCAATGtatggggagacctccactgtggcaagaagattttctttattaatgcatttttaattaca  
gATGGCAAAAACATCATACTCGTTCTACATCGTTCTGAAGGATCAATTTTAAaggggaaactccccac  
ccgaccagacgacggaaagctaacgatgtgcaattgaatagtcatttagtcgttttgcgcgaacgaactaacctttgacttttaagttcactacggt  
gaggacaaaaatcaataaattaaatcgagaccgttgatgagcaaaagaaaaaaaataattttactgattttcatttcgttccatcgactacataatcataattat  
atgccacattttattataagttttg

SEQ ID NO:22

35 Genomic Nucleic Acid Sequence  
931 nucleotides  
Mosquito odorant receptor 6

40 aacacccatcttatcggcaaaattagattttaccgtttgaaagcgggttcccttccctggctgtttctcactctctctctctctctctcttattgatgccgtatgcg  
ccgcgtgctataggctagTTATGCTTACCGGATGTTGCGATCGCGCACGTGCTTTTCCGCATACGCCAG  
TGCACACTTGATGGCGGTGGTGTGACGTCTGCTGCGCACCGTTTTCTGCTCGTGAGTCAGA  
CCTTTTCATTTCTGCAATATCCTGTTTCTTTCCCGACCCACAGACGGTTAGACGGATATAT  
GCTGGTAAAGTTTGTCTCTTCATGCTGTGCTTTCTGATCGAGCTGCTGATGCTGTGTGCGTA  
CGGTGAGGATATTGTGGAATCGgtaaggcaccaggcgggtgatgagcgagtcgcgagtaattgaagcttttgccttttaaacacatca  
45 gagCCTTGGGGTGATTGATGCCGCTTACGGTTGCGAATGGTACCGGGAAGGGTCGGTGGCGTT  
CCATCGATCCGTGCTGCAAATTATACACCGCAGCCAGCAGTCCGTCATACTGACCGCATGGA

AAATTTGGCCCATCCAAATGAGTACTTTCAGTCAGgtgagttgccaattgattgccgtttgcgttaattttcagtaagagt  
gcgctctttcccttagATCCTGCAAGCTTCCTGGTCCTACTTTACCCTCCTGAAGACCGTCTACGGGAA  
TAAgtaagcgcgagagagagagagagagcagtatcgttcaccctttggatgaatcaatagatttctaatacatgaaccattgaaaaatgaatcaacatttt  
cgctagttgcacaatatgtaccattctatacagcttcaccacgaccaagcgtttgttgcacaggaccaaacacgtttcgacaagccgcgtcacctgctgg  
c

**SEQ ID NO:23**  
Genomic Nucleic Acid Sequence  
11,103 nucleotides  
Mosquito odorant receptor 7

ccgccccgggcaggtgacttacgcggtctgacttgctggtgcgctgctttgtacggcaaacggctacacaagcgaatcgaattatttctatcacgctgcg  
cttaccagcgcctgctggtaggcaaagaatgtgcaaagtttcatttggcttggttcgctgctttgctgtgaacgtgtgcacggttgcacgctaaggtttcgg  
tgtgagccgagaagttgcagatcgaatctctttgtgtgtgtgtgtgtgtgcagtggaagcattgtgtttagtgagaagtgaagaaagtgctgaaa  
aatgcaagtcacagccgaccaagtacgtcggccttcgttgcgacgtgatgccgaacattcgggttgatgcaggccagcgggtcaactttctgttccggctac  
gtcaccggccccgatactgacgcgaaggtgtactcctggtggacgctcgcctATGGTGCTGATCCAGTTCTTCGCCATCCTC  
GGCAACCTGGCGACGAACGCGGACGACGTGAACGAGCTGACCGCCAACACGATCACGACCC  
TGTTCTTCACGCACTCGGTCACCAAGTTCATCTACTTTGCGGTCAACTCGGAGAACTTCTACC  
GGACGCTCGCCATCTGGAACCAGACCAACACGCACCCGCTGTTTGCCGAATCGGACGCCCCG  
GTACCATTTCGATTGCGCTCGCCAAGATGCGGAAGCTGCTGGTGCTGGTGATGGCCACCACCG  
TCCTGTCGGTTGTCGgtatgtgtgtatgtgtgtggcgtttgggaaagtgtctttgcggcagaaccccaatctactgttacgcttgactgggtttt  
tgttttttctcgggtggaggacgggataaaatatctgaagaataattgagtcacccacaggggatgcaagacatcgcaggcagagagtttgggttg  
atttatcaccgcacaccgaatatcttcacggttcataagcttcaccgcggtgaaaagggaactccccatttcctgttttttttctcctctcgataaattac  
tcategcttttcgtttttttttgttgttgccttcttcttcttcatccctactagCCTGGGTTACGATAACATTTTTCGGCGAGAGC  
GTCAAGACTGTGCTCGATAAGGCAACCAACGAGACGTACACGGTGGATATACCCCGGCTGC  
CCATCAAGTCCTGGTATCCGTGGAATGCAATGAGCGGACCGGCGTACATTTTCTCTTTCATCT  
ACCAGGTACGTTGGCGGAATgtcctgcgcgtcacagttggcagtcagtgagcggcaacacggcgaaaaaatgggactaaaaccgg  
tcttcacagagccaacacattcctacagcaattgcataccttcgggcggtcgggactgggcaatgcagctacaacatcctcgcctaaagtattgaattcg  
agcgacaaatgttgccgtgttagggcgttttgtgataatagtcgtttttgtcctcgccttatcaaacggaggaaatccatttcgctacaatgcc  
tacagctcaagtttcaaggtcaatcgagcgggtgggatcaacttttttattcattttgctaacgccccatcaacaaattctatgttctcaatggcaagattact  
gcccgcaccaatcgcacaacgaaacggcaaaagaaaagcgacgattatgaagatgtcaaaccattgcccgcgcgacgctttatctgatgatttcggg  
atggcttttacttgctgtactttcaggcacaaaaggaaatgaaaccagcgcaggtcgtttgccggcttgaggaggttcttcaggcactgaggctgagta  
cttaaatcgaacgatttttacgattctggatccagttttatgatgtggcctgcattacagtggaattataacctgatgttcatttcattgcattttgtaagttgtgct  
ggtaacgcccgtaacgattaattctttcaaagagattctttcaaagagattcaaatgtgtataacaaatgctaacgaatggaccgtacttgagggttgcg  
gaaagtaacgttttaaaatattcatcacaatcctctgcaaactgtgcttaattaattggtgcacaataagtttaactgtggcggcagatgtgtcgtgtccgc  
ttccttcttcccagcaagctcgtgcgaataatttattccatcatttaataacagccgtttgtgcattttaattagcaaagcaatataaaaagcagctaaccatc  
cccattaaaacaaagtgttccgggcccattgttatggcggtggaaagtaattggtttaccagtggaggtgtccttcccatcgtgggtacttcgcgatattc  
ttgtcttatacaagtgcatacagaaaaaaggacaaatcctccttgcctatggtctaaaggccagcttcgggtaccgcttccgcttcgggatgtcataaagttgat  
gggtgttttaacattacttcgctttaaaccacctaattggacttttcatgcttgagctaaagttaaaccagccaccagcggtagcaccagccacgggtgat  
ttcggcgggcctcatccccagttttgcgccaccaatattgccttcattaatctgtaccctcggagcgttagggcccgcggacgagtcctcgttgtaatgc  
accgccatgccacgggacgggataatccgttgggacggcgcgaaagcgactatcgcggacggattggttcgaccgtgctacaacacattttatgcttca  
cagatttacttctgctgttttcgatggtccagagcaacctcgcggatgtcatgttctgctcctggttgctgctagcctgcgagcagctgcaacactgaaggt

5

10

15

20

:25

30

35

40

45



20437054907

aggttggtgcttccacacgagcattgctttccgtaccgcggtggggcgagtttcaacgcaaccttctacaagcaacgccacaacgcctgggagcgatat  
ttaaagaaacaagaacatcccgaacttcagcacatgccgtgattgacctgttgaaaagctttgtgagcgtgtgagttgaacgagctctatttccagcg  
atgggtggcatttgtgtggcatgctatcgtcagcttttctgaatctttacctctccattcgcctccattagtagacgcgtatggaaaatgggtgcaacggatca  
gaacggatttccgcgacagacttaataaagggaagcaacgcggttttgcattgtgtagtgtttatgagctttatgccgttactttgcaattaaaaatagcaaa  
5 aaataacagtttttttgaagcggattacaaagaatgtatcagaatattacgtgaaacattcatttcatgctgttaacgctcaaatagattttgtaacacg  
gattgcataaccttgcggtatcggttacatttgcctaacagtatgcaatctgttttagctttgttttaagtactgcgttggttagtacaattttttacaccgc  
gtaatttatctcacaattgcaaaaaatgcaatctgtatcgatttccacacaaatcagatcccgaaccagtgtagcccaatgtgctcttattgaattacca  
cgaacaaatcaacctgatgcccggttccgttggaacagcttgcgcgaagccgctcagtggttctgactaccgtgctgccattttgctgccctcatc  
gaacagataaacagaagggaactcttgtgagcatcgcaatgcccgctgaagtccgctgaaaatgggctaaattcaatttgacgcatttaccgcgaa  
10 caattgcgcgaaggctgtcaagtgtgttccacgaactgcgacaacaagcacacacacacaaatgttatcgttccgcatgtttctcggtacaaagcg  
tgtggcgtatgtggcatgccgattcccagacagagtgatcgatagtaaatgtagcctatccggtagcattcaatttcttctatcctcgaaacaaagcc  
cattctggggaggcggtgtgaagcttcaaggcattgtgaaacaaatgtcctgggtcggagggatgctggggaaagcaaacacgggtgccgccatcgct  
gctaccgtcaatcgatcatgcatgtgtattaattttgtttattccctgcgtatctatgcgtccgctgctgcttccgatttccggaagtcaaggaaaaag  
cgactccatttgggattgttttgcagcgaaaaaatcaaaacattcgcacaaaaccgctctccatttcaaatgcctacactgtcactgtatatctcttctctc  
15 gtttgcacgttgcagTCTCGTTTCAGCAATCGGAGATACGTACGGTCCTGCCCTGCTGCTACACATG  
CTGACCTCCACCATCAAGCTGACGCTGCTCGCCTACCAGGCAACGAAAATCGACGGTGTCAA  
CGTGTACGGATTGACCGTAATCGGATATTTGTGCTACGCGTTGGCTCAGGTTTTCTCTGTTTTG  
CATCTTTGGCAATCGGCTCATCGAGGAGgtacgtgcgtcggcggttgccgtgggaaagcatttccctgccccatctcgtt  
catttcccagatcacacatttgcatacaaaagccagcacacttttgcctcgcgctgccatctcggttctgaattgtttcacttctcccatacttctcccgtgc  
20 agAGCTCATCCGTGATGAAGGCGGCCTATTCCTGCCACTGGTACGACGGGTCCGAGGAGGCA  
AAAACCTTCGTCCAGATCGTTTGTGTCAGCAGTGCCAGAAGGCGATGACTATTTCCGGAGCCAA  
GTTTTTCACCGTTTCGCTCGATCTGTTTGCTTCGtaagtgtagcctgggtggctggcacagaacagggtggcaaaacagg  
gactttggctctagcctgatgggtggtatgtgtgtctatttttgcctaccattctcgcaccccttcttccagGTTCTTGGAGCCGTTGTCA  
CCTACTTCATGGTGCTGGTGCAGCTGAAGTAAacagccgtggcccgaaggatgtgttttttgcctcgttccggttgttgtt  
25 gtgcacacttctcttggacatttctctactgcaaagggttaacaaacagcaacaacaaataatcccaagtttctttacagatctttgcaaaatgattagattt  
aatagattaacagtgttgattatctgtcctgtagcaaccggggctgaagaacgttgatttgtaaaagtacaaaaggacgttggaattgaaccaccag  
aagagtataatttatgcaaaagtcaccaagggaatctatgtatgtgtgatttgcgtcatcaagcactgtatgtgcctttcaactagtgcagcaataaagagt  
acaaatgtttcttagcgcaccgtacattgtcgttccggcggtttaaaccgttgtgataatacacaacaaatgataaaaaataataacaaatgttaatatgag  
taagtactaaatagagaaatcgttttagtatgatcatacctccaatcattgtttgaaattaactttaatttaactcaaaatgaaccgatgtttacttctgtgagaa  
30 ttattgtggaagaacttaattggaagtataattaaattgattgctaactttatgcgttttcaatttacgaacgctagtcttcaaacatcgcttcaaaagtattactac  
cacattattcatttactttagttatatttgcctcttcatcttccatggccagaactactgcagaaaagcttcttttgcctcgttccgatggttggttgacg  
aagttggtacaaaacggcaagcaattagcataaactatttgcacgcagatggaaatgaatgtaccactagaaccgagtgaatgaattactttcaacttg  
cacgcaaaaaccattatctaaagtacgcacaacttaaaaacaaaccccaattgtcgtccacccttccattccatttctgtacactttccgaccgagttctgt  
agcgcacagcagcaaaaaatacatataaaaccttcatcactcaagctgtatcgagccagcgtgggttgtgtttgactgtgctgtgaaagaaagaagaaaa  
35 aaaaaacacttccacgggaagctagcaattggaatgcataaattaaccggaagaaatcgcaaaaccccgaccgacgtaccgcaccgcacccgtac  
cgataccggaacaaacgggtgtgcgcgaaagaatccgctagcagccccactggcacgggtatttgcgttctgtgttttcttccactggttgggtgcc  
tgggcgaaggctagctcggctacttcccgggcgcaatttctgcagcccaaggcggtgctcgtggggccaaaagaat